

THE REVERSE YANG-MILLS-HIGGS FLOW IN A NEIGHBOURHOOD OF A CRITICAL POINT

GRAEME WILKIN

ABSTRACT. The main result of this paper is a construction of solutions to the reverse Yang-Mills-Higgs flow converging in the C^∞ topology to a critical point. The construction uses only the complex gauge group action, which leads to an algebraic classification of the isomorphism classes of points in the unstable set of a critical point in terms of a filtration of the underlying Higgs bundle.

Analysing the compatibility of this filtration with the Harder-Narasimhan-Seshadri double filtration gives an algebraic criterion for two critical points to be connected by a flow line. As an application, we can use this to construct Hecke modifications of Higgs bundles via the Yang-Mills-Higgs flow. When the Higgs field is zero (corresponding to the Yang-Mills flow), this criterion has a geometric interpretation in terms of secant varieties of the projectivisation of the underlying bundle inside the unstable manifold of a critical point, which gives a precise description of broken and unbroken flow lines connecting two critical points. For non-zero Higgs field, at generic critical points the analogous interpretation involves the secant varieties of the spectral curve of the Higgs bundle.

1. INTRODUCTION

There is a well-known relationship between the Yang-Mills heat flow on a Riemann surface and the notion of stability from algebraic geometry. This began with work of Atiyah and Bott [1] and continued with Donaldson's proof [7] of the Narasimhan-Seshadri theorem [39] and subsequent work of Daskalopoulos [5] and Rade [41], which shows that the Yang-Mills flow converges to a unique critical point which is isomorphic to the graded object of the Harder-Narasimhan-Seshadri double filtration of the initial condition. In the setting of Higgs bundles, a theorem of Hitchin [15] and Simpson [45] shows that a polystable Higgs bundle is gauge equivalent to the minimum of the Yang-Mills-Higgs functional and that this minimum is achieved by the heat flow on the space of metrics. The results of [51] show that the theorem of Daskalopoulos and Rade described above extends to the Yang-Mills-Higgs flow on the space of Higgs bundles over a compact Riemann surface. More generally, when the base manifold is compact and Kähler, then these results are due to [8], [9], [49], [45], [6], [43], [22] and [32].

Continuing on from these results, it is natural to investigate flow lines between critical points. Naito, Kozono and Maeda [34] proved the existence of an unstable manifold of a critical point for the Yang-Mills functional, however their method does not give information about the isomorphism classes in the unstable manifold, and their proof requires a manifold structure on the space of connections (which is not true for the space of Higgs bundles). Recent results of Swoboda [48] and Janner-Swoboda [23] count flow lines for a perturbed Yang-Mills functional, however these perturbations destroy the algebraic structure of the

Date: March 29, 2018.

This research was supported by grant number R-146-000-200-112 from the National University of Singapore. The author also acknowledges support from NSF grants DMS 1107452, 1107263, 1107367 "RNMS GEometric structures And Representation varieties" (the GEAR Network).

Yang-Mills flow, and so there does not yet exist an algebro-geometric description of the flow lines in the spirit of the results described in the previous paragraph. Moreover, one would also like to study flow lines for the Yang-Mills-Higgs functional, in which case the perturbations do not necessarily preserve the space of Higgs bundles, which is singular.

The purpose of this paper is to show that in fact there is an algebro-geometric description of the flow lines connecting given critical points of the Yang-Mills-Higgs functional over a compact Riemann surface. As an application, we show that the Hecke correspondence for Higgs bundles studied by Witten in [52] has a natural interpretation in terms of gradient flow lines. Moreover, for the Yang-Mills flow, at a generic critical point there is a natural embedding of the projectivisation of the underlying bundle inside the unstable set of the critical point, and the results of this paper show that the isomorphism class of the limit of the downwards flow is determined if the initial condition lies in one of the secant varieties of this embedding, giving us a geometric criterion to distinguish between broken and unbroken flow lines. For the Yang-Mills-Higgs flow the analogous picture involves the secant varieties of the space of Hecke modifications compatible with the Higgs field. At generic critical points of the Yang-Mills-Higgs functional this space of Hecke modifications is the spectral curve of the Higgs bundle.

The basic setup for the paper is as follows. Let $E \rightarrow X$ be a smooth complex vector bundle over a compact Riemann surface with a fixed Hermitian metric and let \mathcal{B} denote the space of Higgs pairs on E . The *Yang-Mills-Higgs functional* is

$$\text{YMH}(\bar{\partial}_A, \phi) := \|F_A + [\phi, \phi^*]\|_{L^2}^2$$

and the *Yang-Mills-Higgs flow* is the downwards gradient flow of YMH given by the equation (2.4). This flow is generated by the action of the complex gauge group $\mathcal{G}^{\mathbb{C}}$. Equivalently, one can fix a Higgs pair and allow the Hermitian metric on the bundle to vary in which case the flow becomes a nonlinear heat equation on the space of Hermitian metrics (cf. [8], [45]). At a critical point for this flow the Higgs bundle splits into Higgs subbundles and on each subbundle the Higgs structure minimises YMH. The *unstable set* of a critical point $(\bar{\partial}_A, \phi)$ consists of all Higgs pairs for which a solution to the YMH flow (2.4) exists for all negative time and converges in the smooth topology to $(\bar{\partial}_A, \phi)$ as $t \rightarrow -\infty$. The first theorem of the paper gives an algebraic criterion for a complex gauge orbit to intersect the unstable set for the Yang-Mills-Higgs flow.

Theorem 1.1 (Criterion for convergence of reverse heat flow). *Let E be a complex vector bundle over a compact Riemann surface X , and let $(\bar{\partial}_A, \phi)$ be a Higgs bundle on E . Suppose that E admits a filtration $(E^{(1)}, \phi^{(1)}) \subset \dots \subset (E^{(n)}, \phi^{(n)}) = (E, \phi)$ by Higgs subbundles such that the quotients $(Q_k, \phi_k) := (E^{(k)}, \phi^{(k)}) / (E^{(k-1)}, \phi^{(k-1)})$ are Higgs polystable and $\text{slope}(Q_k) < \text{slope}(Q_j)$ for all $k < j$. Then there exists $g \in \mathcal{G}^{\mathbb{C}}$ and a solution to the reverse Yang-Mills-Higgs heat flow equation with initial condition $g \cdot (\bar{\partial}_A, \phi)$ which converges to a critical point isomorphic to $(Q_1, \phi_1) \oplus \dots \oplus (Q_n, \phi_n)$.*

Conversely, if there exists a solution of the reverse heat flow from the initial condition $(\bar{\partial}_A, \phi)$ converging to a critical point $(Q_1, \phi_1) \oplus \dots \oplus (Q_n, \phi_n)$ where each (Q_j, ϕ_j) is polystable with $\text{slope}(Q_k) < \text{slope}(Q_j)$ for all $k < j$, then (E, ϕ) admits a filtration $(E^{(1)}, \phi^{(1)}) \subset \dots \subset (E^{(n)}, \phi^{(n)}) = (E, \phi)$ whose graded object is isomorphic to $(Q_1, \phi_1) \oplus \dots \oplus (Q_n, \phi_n)$.

A key difficulty in the construction is the fact that the space of Higgs bundles is singular, and so the existing techniques for constructing unstable sets (see for example [34] for the Yang-Mills flow or [24, Sec. 6] in finite dimensions) cannot be directly applied since they depend on the manifold structure of the ambient space. One possibility is to study the unstable set of the function $\|F_A + [\phi, \phi^*]\|_{L^2}^2 + \|\bar{\partial}_A \phi\|_{L^2}^2$ on the space of all pairs $(\bar{\partial}_A, \phi)$ without the Higgs bundle condition $\bar{\partial}_A \phi = 0$, however one would then need a criterion to determine when a point in this unstable set is a Higgs bundle and one would also need a method to determine the isomorphism classes of these points.

The construction in the proof of Theorem 1.1 is intrinsic to the singular space since it uses the action of the complex gauge group to map the unstable set for the linearised YMH flow (for which we can explicitly describe the isomorphism classes) to the unstable set for the Yang-Mills-Higgs flow. The method used here to compare the flow with its linearisation is called the “scattering construction” in [19] and [40] since it originates in the study of wave operators in quantum mechanics (see [42] for an overview). The method in this paper differs from [19] and [40] in that (a) the construction here is done using the gauge group action in order to preserve the singular space and (b) the distance-decreasing formula for the flow on the space of metrics [8] is used here in order to avoid constructing explicit local coordinates as in [19] (the construction of [19] requires a manifold structure around the critical points).

As a consequence of Theorem 1.1, we have an algebraic criterion for critical points to be connected by flow lines.

Corollary 1.2 (Algebraic classification of flow lines). *Let $x_u = (\bar{\partial}_{A_u}, \phi_u)$ and $x_\ell = (\bar{\partial}_{A_\ell}, \phi_\ell)$ be critical points with $\text{YMH}(x_u) > \text{YMH}(x_\ell)$. Then x_u and x_ℓ are connected by a flow line if and only if there exists a Higgs pair (E, ϕ) which has Harder-Narasimhan-Seshadri double filtration whose graded object is isomorphic to x_ℓ , and which also admits a filtration $(E^{(1)}, \phi^{(1)}) \subset \dots \subset (E^{(n)}, \phi^{(n)}) = (E, \phi)$ by Higgs subbundles such that the quotients $(Q_k, \phi_k) := (E^{(k)}, \phi^{(k)}) / (E^{(k-1)}, \phi^{(k-1)})$ are polystable and satisfy $\text{slope}(Q_k) < \text{slope}(Q_j)$ for all $k < j$ and the graded object $(Q_1, \phi_1) \oplus \dots \oplus (Q_n, \phi_n)$ is isomorphic to x_u .*

As an application of the previous theorem, we can construct Hecke modifications of Higgs bundles via Yang-Mills-Higgs flow lines. First consider the case of a Hecke modification at a single point (miniscule Hecke modifications in the terminology of [52]).


Theorem 1.3. (1) *Let $0 \rightarrow (E', \phi') \rightarrow (E, \phi) \xrightarrow{v} \mathbb{C}_p \rightarrow 0$ be a Hecke modification such that (E, ϕ) is stable and (E', ϕ') is semistable, and let L_u be a line bundle with $\deg L_u + 1 < \text{slope}(E') < \text{slope}(E)$. Then there exist sections $\phi_u, \phi_\ell \in H^0(K)$, a line bundle L_ℓ with $\deg L_\ell = \deg L_u + 1$ and a metric on $E \oplus L_u$ such that $x_u = (E, \phi) \oplus (L_u, \phi_u)$ and $x_\ell = (E'_{gr}, \phi'_{gr}) \oplus (L_\ell, \phi_\ell)$ are critical points connected by a YMH flow line, where (E'_{gr}, ϕ'_{gr}) is isomorphic to the graded object of the Seshadri filtration of (E', ϕ') .*

(2) *Let $x_u = (E, \phi) \oplus (L_u, \phi_u)$ and $x_\ell = (E', \phi') \oplus (L_\ell, \phi_\ell)$ be critical points connected by a YMH flow line such that L_u, L_ℓ are line bundles with $\deg L_\ell = \deg L_u + 1$, (E, ϕ) is stable and (E', ϕ') is polystable with $\deg L_u + 1 < \text{slope}(E') < \text{slope}(E)$. Then (E', ϕ') is the graded object of the Seshadri filtration of a Hecke modification of (E, ϕ) . If (E', ϕ') is Higgs stable then it is a Hecke modification of (E, ϕ) .*


For Hecke modifications defined at multiple points, we can inductively apply the above theorem to obtain a criterion for two critical points to be connected by a broken flow line. For non-negative integers m, n , the definition of (m, n) stability is given in Definition 4.5. The space $\mathcal{N}_{\phi, \phi_u}$ denotes the space of Hecke modifications compatible with the Higgs fields ϕ and ϕ_u (see Definition 4.11).

Corollary 1.4. *Consider a Hecke modification $0 \rightarrow (E', \phi') \rightarrow (E, \phi) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$ defined by $n > 1$ distinct points $\{v_1, \dots, v_n\} \in \mathbb{P}E^*$, where (E, ϕ) is $(0, n)$ stable. If there exists $\phi_u \in H^0(K)$ such that $v_1, \dots, v_n \in \mathcal{N}_{\phi, \phi_u}$, then there is a broken flow line connecting $x_u = (E, \phi) \oplus (L_u, \phi_u)$ and $x_\ell = (E'_{gr}, \phi'_{gr}) \oplus (L_\ell, \phi_\ell)$, where (E'_{gr}, ϕ'_{gr}) is the graded object of the Seshadri filtration of the semistable Higgs bundle (E', ϕ') .*

For any gradient flow, given upper and lower critical sets C_u and C_ℓ , one can define the spaces $\mathcal{F}_{\ell, u}$ (resp. $\mathcal{BF}_{\ell, u}$) of unbroken flow lines (resp. broken or unbroken flow lines) connecting these sets, and the spaces $\mathcal{P}_{\ell, u}$ (resp. $\mathcal{BP}_{\ell, u}$) of pairs of critical points connected by an unbroken flow line (resp. broken or unbroken flow line). These spaces are correspondences with canonical projection maps to the critical sets given by the projection taking a flow line to its upper and lower endpoints.



In the setting of Theorem 1.3, let $d = \deg E$ and $r = \text{rank}(E)$ and let C_u and C_ℓ be the upper and lower critical sets. There are natural projection maps to the moduli space of semistable Higgs bundles $C_u \rightarrow \mathcal{M}_{ss}^{Higgs}(r, d)$ and $C_\ell \rightarrow \mathcal{M}_{ss}^{Higgs}(r, d-1)$. Suppose that $\gcd(r, d) = 1$ so that $\mathcal{M}_{ss}^{Higgs}(r, d)$ consists solely of stable Higgs pairs and hence any Hecke modification is semistable. Since the flow is \mathcal{G} -equivariant, then there is an induced correspondence variety, denoted $\mathcal{M}_{\ell, u}$ in the diagram below.



As a consequence of Theorem 1.3, we have the following result.

Corollary 1.5. *$\mathcal{M}_{\ell, u}$ is the Hecke correspondence.*

A natural question motivated by Floer theory is to ask whether a pair of critical points connected by a broken flow line can also be connected by an unbroken flow line, i.e whether $\mathcal{BP}_{\ell,u} = \mathcal{P}_{\ell,u}$. For example, in the context of the Yang-Mills flow, Swoboda [48] constructs a Morse complex by counting flow lines between critical points of a perturbed Yang-Mills functional. The appearance of the Hecke correspondence in previous theorem shows that spaces of flow lines for the *unperturbed* functional carry a lot of interesting algebraic structure and a natural next step is to develop a criterion distinguishing between broken and unbroken flow lines. The methods used to prove the Theorem 1.3 and Corollaries 1.4 and 1.5 can be used to investigate this question using the geometry of secant varieties of the space of Hecke modifications inside the unstable set of a critical point. For critical points of the type studied in Theorem 1.3, generically this space of Hecke modifications is the spectral curve of the Higgs field, and so the problem reduces to studying secant varieties of the spectral curve. This is explained in detail in Section 4.4. In particular, Corollary 4.27 gives a complete classification of the unbroken flow lines on the space of rank 2 Higgs bundles.

Another direction is to apply the methods of this paper to other geometric contexts, such as the Yang-Mills flow on higher-dimensional compact Kähler manifolds where bubbling occurs (cf. [6], [22], [43]). For a perturbed heat flow on the loop space of a compact Riemannian manifold, Weber [50] has proved a backward λ -lemma around a non-degenerate critical point, and it would be interesting to use the techniques here to obtain information about flow lines for the unperturbed flow. Another case of interest is the nonlinear Cauchy-Riemann equations studied in Floer theory [11], [12], [13]. The bounded trajectories studied in [13] have many properties in common with spaces of flow lines in finite-dimensional Morse theory, and it would also be interesting to see if the methods used here also apply in the setting of [13].

The paper is organised as follows. In Section 2 we set the notation for the paper, prove a slice theorem around the critical points and derive some preliminary estimates for the YMH flow near a critical point. Section 3 contains the main part of the analysis of the YMH flow around a critical point, which leads to the proof of Theorem 1.1 and Corollary 1.2. In Section 4 we interpret the analytic results on flow lines in terms of the Hecke correspondence, leading to the proof of Theorem 1.3, Corollary 1.4 and Corollary 1.5. Appendix A contains a proof that a solution to the reverse YMH flow with a given initial condition is necessarily unique.

Acknowledgements. I would like to thank George Daskalopoulos, Mudumbai Narasimhan and Richard Wentworth for their interest in the project, as well as George Hitching for useful discussions about [4] and [17].

2. PRELIMINARIES

2.1. The Yang-Mills-Higgs flow on a compact Riemann surface. Fix a compact Riemann surface X and a smooth complex vector bundle $E \rightarrow X$. Choose a normalisation so that $\text{vol}(X) = 2\pi$. Fix $\bar{\partial}_{A_0} : \Omega^0(E) \rightarrow \Omega^{0,1}(E)$ such that $\bar{\partial}_{A_0}$ is \mathbb{C} -linear and satisfies the Leibniz rule $\bar{\partial}_{A_0}(fs) = (\bar{\partial}f)s + f(\bar{\partial}_{A_0}s)$ for all $f \in \Omega^0(X)$ and $s \in \Omega^0(E)$. Let $\mathcal{A}^{0,1}$ denote the affine space $\bar{\partial}_{A_0} + \Omega^{0,1}(\text{End}(E))$. A theorem of Newlander and Nirenberg identifies $\mathcal{A}^{0,1}$ with the space of holomorphic structures on E . The *space of Higgs bundles on E* is

$$(2.1) \quad \mathcal{B} := \{(\bar{\partial}_A, \phi) \in \mathcal{A}^{0,1} \times \Omega^{1,0}(\text{End}(E)) : \bar{\partial}_A \phi = 0\}$$

The complex gauge group is denoted $\mathcal{G}^{\mathbb{C}}$ and acts on \mathcal{B} by $g \cdot (\bar{\partial}_A, \phi) = (g\bar{\partial}_A g^{-1}, g\phi g^{-1})$. If X is a complex manifold with $\dim_{\mathbb{C}} X > 1$ then we impose the extra integrability conditions $(\bar{\partial}_A)^2 = 0$ and $\phi \wedge \phi = 0$. Given a Hermitian metric on E , let \mathcal{A} denote the space of connections on E compatible with the metric, and let $\mathcal{G} \subset \mathcal{G}^{\mathbb{C}}$ denote the subgroup of unitary gauge transformations. The Chern connection construction defines an injective map $\mathcal{A}^{0,1} \hookrightarrow \mathcal{A}$ which is a diffeomorphism when $\dim_{\mathbb{C}} X = 1$. Given $\bar{\partial}_A \in \mathcal{A}^{0,1}$, let F_A denote the curvature of the Chern connection associated to $\bar{\partial}_A$ via the Hermitian metric. The metric induces a pointwise norm $|\cdot| : \Omega^2(\text{End}(E)) \rightarrow \Omega^0(X, \mathbb{R})$ and together with the Riemannian structure on X an L^2 norm $\|\cdot\|_{L^2} : \Omega^2(\text{End}(E)) \rightarrow \mathbb{R}$. The *Yang-Mills-Higgs functional* $\text{YMH} : \mathcal{B} \rightarrow \mathbb{R}$ is defined by

$$(2.2) \quad \text{YMH}(\bar{\partial}_A, \phi) = \|F_A + [\phi, \phi^*]\|_{L^2}^2 = \int_X |F_A + [\phi, \phi^*]|^2 d\text{vol}$$

When $\dim_{\mathbb{C}} X = 1$, the Hodge star defines an isometry $*$: $\Omega^2(\text{End}(E)) \rightarrow \Omega^0(\text{End}(E)) \cong \text{Lie } \mathcal{G}^{\mathbb{C}}$. For any initial condition (A_0, ϕ_0) , the following equation for $g_t \in \mathcal{G}^{\mathbb{C}}$ has a unique solution on the interval $t \in [0, \infty)$ (cf. [8], [45])

$$(2.3) \quad \frac{\partial g}{\partial t} g_t^{-1} = -i * (F_{g_t \cdot A_0} + [g_t \cdot \phi_0, (g_t \cdot \phi_0)^*]), \quad g_0 = \text{id}.$$

This defines a unique curve $(A_t, \phi_t) = g_t \cdot (A_0, \phi_0) \in \mathcal{B}$ which is a solution to the downwards Yang-Mills-Higgs gradient flow equations

$$(2.4) \quad \begin{aligned} \frac{\partial A}{\partial t} &= i\bar{\partial}_A * (F_A + [\phi, \phi^*]) \\ \frac{\partial \phi}{\partial t} &= i[\phi, *(F_A + [\phi, \phi^*])]. \end{aligned}$$

for all $t \in [0, \infty)$. The result of [51, Thm 3.1] shows that the solutions converge to a unique limit $(A_{\infty}, \phi_{\infty})$ which is a critical point of YMH. Moreover [51, Thm. 4.1] shows that the isomorphism class of this limit is determined by the graded object of the Harder-Narasimhan-Seshadri double filtration of the initial condition (A_0, ϕ_0) .

Remark 2.1. Since the space \mathcal{B} of Higgs bundles is singular, then we define the gradient of YMH as the gradient of the function $\|F_A + [\phi, \phi^*]\|_{L^2}^2$ defined on the ambient smooth space $T^*\mathcal{A}^{0,1}$, which contains the space \mathcal{B} as a singular subset. When the initial condition is a Higgs bundle, then a solution to (2.4) is generated by the action of the complex gauge group $\mathcal{G}^{\mathbb{C}}$ which preserves \mathcal{B} . Therefore the solution to (2.4) is contained in \mathcal{B} and so from now on we can consider the flow (2.4) as a well-defined gradient flow on the singular space \mathcal{B} . Throughout the paper we define a critical point to be a stationary point for the Yang-Mills-Higgs flow.

Definition 2.2. A *critical point* for YMH is a pair $(A, \phi) \in \mathcal{B}$ such that

$$(2.5) \quad \bar{\partial}_A * (F_A + [\phi, \phi^*]) = 0, \quad \text{and} \quad [\phi, *(F_A + [\phi, \phi^*])] = 0.$$

The critical point equations (2.5) imply that the bundle E splits into holomorphic ϕ -invariant sub-bundles $E_1 \oplus \cdots \oplus E_n$, such that the induced Higgs structure $(\bar{\partial}_{A_j}, \phi_j)$ on the bundle E_j minimises the Yang-Mills-Higgs functional on the bundle E_j (cf. [1, Sec. 5] for holomorphic bundles and [51, Sec. 4] for Higgs bundles). In particular, each Higgs pair $(\bar{\partial}_{A_j}, \phi_j)$ is polystable. The decomposition is not necessarily

unique due to the possibility of polystable bundles with the same slope, however it is unique if we impose the condition that $(E_1, \phi_1) \oplus \cdots \oplus (E_n, \phi_n)$ is the graded object of the socle filtration of the Higgs bundle (E, ϕ) (see [20] for holomorphic bundles and [2, Sec. 4] for Higgs bundles). With respect to this decomposition the curvature $*(F_A + [\phi, \phi^*]) \in \Omega^0(\text{ad}(E)) \cong \text{Lie}(\mathcal{G})$ has the following block-diagonal form

$$(2.6) \quad i * (F_A + [\phi, \phi^*]) = \begin{pmatrix} \nu_1 \cdot \text{id}_{E_1} & 0 & \cdots & 0 \\ 0 & \nu_2 \cdot \text{id}_{E_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \nu_n \cdot \text{id}_{E_n} \end{pmatrix}$$

where $\nu_j = \text{slope}(E_j)$ and we order the eigenvalues by $\nu_j < \nu_k$ for all $j < k$.

Definition 2.3. A *Yang-Mills-Higgs flow line* connecting an upper critical point $x_u = (\bar{\partial}_{A_u}, \phi_u)$ and a lower critical point $x_\ell = (\bar{\partial}_{A_\ell}, \phi_\ell)$ is a continuous map $\gamma : \mathbb{R} \rightarrow \mathcal{B}$ such that

- (1) $\frac{d\gamma}{dt}$ satisfies the Yang-Mills-Higgs flow equations (2.4), and
- (2) $\lim_{t \rightarrow -\infty} \gamma(t) = x_u$ and $\lim_{t \rightarrow \infty} \gamma(t) = x_\ell$, where the convergence is in the C^∞ topology on \mathcal{B} .

Definition 2.4. The *unstable set* $W_{x_u}^-$ of a non-minimal critical point $x_u = (\bar{\partial}_{A_u}, \phi_u)$ is defined as the set of all points $y_0 \in \mathcal{B}$ such that a solution y_t to the Yang-Mills-Higgs flow equations (2.4) exists for all $(-\infty, 0]$ and $y_t \rightarrow x$ in the C^∞ topology on \mathcal{B} as $t \rightarrow -\infty$.

2.2. A local slice theorem. In this section we define local slices around the critical points and describe the isomorphism classes in the negative slice.

Definition 2.5. Let $x = (\bar{\partial}_A, \phi) \in \mathcal{B}$. The *slice* through x is the set of deformations orthogonal to the $\mathcal{G}^{\mathbb{C}}$ orbit at x .

$$(2.7) \quad S_x = \{(a, \varphi) \in \Omega^{0,1}(\text{End}(E)) \oplus \Omega^{1,0}(\text{End}(E)) \mid \bar{\partial}_A^* a - *[\phi^*, \varphi] = 0, (\bar{\partial}_A + a, \phi + \varphi) \in \mathcal{B}\}.$$

If x is a critical point of YMH with $\beta = *(F_A + [\phi, \phi^*])$, then the *negative slice* S_x^- is the subset

$$(2.8) \quad S_x^- = \{(a, \varphi) \in S_x \mid \lim_{t \rightarrow \infty} e^{i\beta t} \cdot (a, \varphi) = 0\}.$$

Remark 2.6. Near the critical point $x = (\bar{\partial}_A, \phi)$, the dominant term in the gradient flow equation (2.3) for $\frac{\partial g_t}{\partial t} g_t^{-1}$ is $-i*(F_A + [\phi, \phi^*]) = -i\beta$. Therefore the linearised flow with initial condition $x + \delta x$ for $\delta x \in S_x$ has the form $e^{-i\beta t} \cdot (x + \delta x) = x + e^{-i\beta t} \cdot \delta x$, and the negative slice S_x^- is the subset of initial conditions for which the linearised flow converges to the critical point x as $t \rightarrow -\infty$. The action of this linearised flow on $\delta x = (a, \varphi) \in S_x$ is given by conjugation

$$e^{-i\beta t} \cdot (a, \varphi) = \left(e^{-i\beta t} a e^{i\beta t}, e^{-i\beta t} \varphi e^{i\beta t} \right).$$

Therefore, if $i\beta = i*(F_A + [\phi, \phi^*])$ has the block diagonal form of (2.6), then the negative slice consists of all the endomorphisms which are strictly upper-triangular with respect to (2.6).

To prove Lemma 2.8 and Proposition 2.9 below, one needs to first define the slice on the L_1^2 completion of the space of Higgs bundles with the action of the L_2^2 completion of the gauge group. The following lemma shows that if the critical point x is C^∞ then the elements in the slice S_x are also C^∞ .

Lemma 2.7. *Let $x = (\bar{\partial}_A, \phi)$ be a critical point of YMH in the space of C^∞ Higgs bundles, let S_x be the set of solutions to the slice equations in the L_1^2 completion of the space of Higgs bundles and let $\delta x = (a, \varphi) \in S_x$. Then δx is C^∞ .*

Proof. The slice equations are

$$\begin{aligned}\bar{\partial}_A \varphi + [a, \phi] + [a, \varphi] &= 0 \\ \bar{\partial}_A^* a - *[\phi^*, * \varphi] &= 0\end{aligned}$$

Since $(a, \varphi) \in L_1^2$ and $(\bar{\partial}_A, \phi)$ is C^∞ , then the second equation above implies that $\bar{\partial}_A^* a \in L_1^2$ and so $a \in L_2^2$ by elliptic regularity. After applying Sobolev multiplication $L_2^2 \times L_1^2 \rightarrow L^4$, then $[a, \varphi] \in L^4$ and so the first equation above implies that $\bar{\partial}_A \varphi \in L^4$, hence $\varphi \in L_1^4$. Repeating this again shows that $\varphi \in L_2^2$, and then one can repeat the process inductively to show that $\delta x = (a, \varphi)$ is C^∞ . \square

The following result gives a local description of the space of Higgs bundles in terms of the slice. The infinitesimal action of $\mathcal{G}^\mathbb{C}$ at $x \in \mathcal{B}$ is denoted by $\rho_x : \text{Lie}(\mathcal{G}^\mathbb{C}) \cong \Omega^0(\text{End}(E)) \rightarrow \Omega^{0,1}(\text{End}(E)) \oplus \Omega^{1,0}(\text{End}(E))$. Explicitly, for $x = (\bar{\partial}_A, \phi)$ and $u \in \Omega^0(\text{End}(E))$, we have $\rho_x(u) = -(\bar{\partial}_A u, [\phi, u])$. The L^2 -orthogonal complement of $\ker \rho_x \subseteq \Omega^0(\text{End}(E))$ is denoted $(\ker \rho_x)^\perp$.

Lemma 2.8. *Fix $x \in \mathcal{B}$. Then the map $\psi : (\ker \rho_x)^\perp \times S_x \rightarrow \mathcal{B}$ given by $\psi(u, \delta x) = \exp(u) \cdot (x + \delta x)$ is a local homeomorphism.*

Proof. The result of [51, Prop. 4.12] shows that the statement is true for the L_1^2 completion of the space of Higgs bundles and the L_2^2 completion of the gauge group, and so it only remains to show that it remains true on restricting to the space of C^∞ Higgs bundles with the action of the group of C^∞ gauge transformations. The proof of this statement follows from elliptic regularity using the same method as [51, Cor. 4.17]. \square

Now let $x = (\bar{\partial}_A, \phi)$ be a critical point and let $\beta = \mu(x) := *(F_A + [\phi, \phi^*])$. The Lie algebra $\text{Lie}(\mathcal{G}^\mathbb{C}) \cong \Omega^0(\text{End}(E))$ decomposes into eigenbundles for the adjoint action of $e^{i\beta}$. We denote the positive, zero and negative eigenspaces respectively by $\Omega^0(\text{End}(E)_+)$, $\Omega^0(\text{End}(E)_0)$ and $\Omega^0(\text{End}(E)_-)$. The positive and negative eigenspaces are nilpotent Lie algebras with associated unipotent groups $\mathcal{G}_+^\mathbb{C}$ and $\mathcal{G}_-^\mathbb{C}$. The subgroups of \mathcal{G} and $\mathcal{G}^\mathbb{C}$ consisting of elements commuting with $e^{i\beta}$ are denoted \mathcal{G}_β and $\mathcal{G}_\beta^\mathbb{C}$ respectively. Since $\Omega^0(\text{End}(E)_0) \oplus \Omega^0(\text{End}(E)_+)$ is also a Lie algebra then there is a corresponding subgroup denoted $\mathcal{G}_*^\mathbb{C}$.

Let \mathcal{G}_x and $\mathcal{G}_x^\mathbb{C}$ denote the respective isotropy groups of x in \mathcal{G} and $\mathcal{G}^\mathbb{C}$. There is an inclusion $(\mathcal{G}_x)^\mathbb{C} \subseteq \mathcal{G}_x^\mathbb{C}$, however at a non-minimal critical point the two groups may not be equal (in the context of reductive group actions on finite-dimensional affine spaces, this question has been studied by Sjamaar in [47, Prop. 1.6]). At a general critical point, the Higgs bundle (E, ϕ) splits into polystable Higgs sub-bundles $(E_1, \phi_1) \oplus \cdots \oplus (E_n, \phi_n)$, where we order by increasing slope. Then a homomorphism $u \in \text{Hom}(E_j, E_k)$ satisfying $u\phi_j = \phi_k u$ will be zero if $j > k$ since (E_j, ϕ_j) and (E_k, ϕ_k) are polystable and $\text{slope}(E_j) > \text{slope}(E_k)$, however if $j < k$ then the homomorphisms do not necessarily vanish in which case $(\mathcal{G}_x)^\mathbb{C} \subsetneq \mathcal{G}_x^\mathbb{C}$. Therefore $\ker \rho_x = \text{Lie}(\mathcal{G}_x^\mathbb{C}) \subset \Omega^0(\text{End}(E)_+) \oplus \Omega^0(\text{End}(E)_0)$, and so $\mathcal{G}_x^\mathbb{C} \subset \mathcal{G}_*^\mathbb{C}$.

The result of [5, Thm. 2.16] shows that the L_2^2 completion of the gauge group satisfies $\mathcal{G}^{\mathbb{C}} \cong \mathcal{G}_*^{\mathbb{C}} \times_{\mathcal{G}_\beta} \mathcal{G}$. We will use $(\ker \rho_x)_*^\perp$ to denote $(\ker \rho_x)^\perp \cap (\Omega^0(\text{End}(E))_+ \oplus \Omega^0(\text{End}(E)_0))$. At a critical point x , the above argument shows that isotropy group $\mathcal{G}_x^{\mathbb{C}}$ is contained in $\mathcal{G}_*^{\mathbb{C}}$, and so we have the following refinement of Lemma 2.8.

Proposition 2.9. *Let $x \in \mathcal{B}$ be a critical point of YMH. Then there exists a \mathcal{G} -invariant neighbourhood U of x and a neighbourhood U' of $[\text{id}, 0, 0]$ in $\mathcal{G} \times_{\mathcal{G}_\beta} ((\ker \rho_x)_*^\perp \times S_x)$ such that $\psi : U' \rightarrow U$ is a \mathcal{G} -equivariant homeomorphism.*

The results of Section 3 show that the negative slice S_x^- is complex gauge-equivalent to the unstable set W_x^- of a critical point. The next lemma gives a complete classification of the isomorphism classes in S_x^- . Together with the results of Section 3, this is used in Section 4 to classify critical points connected by flow lines.

Lemma 2.10. *Let $x = (E_1, \phi_1) \oplus \cdots \oplus (E_n, \phi_n)$ be a critical point of YMH with curvature as in (2.6) with the Higgs polystable subbundles ordered so that $\text{slope}(E_j) < \text{slope}(E_k)$ iff $j < k$. If $\delta x \in S_x^- \cap U$ then $x + \delta x$ has a filtration $(E^{(1)}, \phi^{(1)}) \subset \cdots \subset (E^{(n)}, \phi^{(n)})$ by Higgs subbundles such that the successive quotients are $(E^{(k)}, \phi^{(k)}) / (E^{(k-1)}, \phi^{(k-1)}) = (E_k, \phi_k)$. Conversely, there exists a neighbourhood U of x such that if a Higgs bundle $y = (E, \phi) \in U$ admits such a filtration then it is gauge equivalent to $x + \delta x$ for some $\delta x \in S_x^-$.*

Proof. The first statement follows directly from the definition of the negative slice in (2.8).

Let $\text{End}(E)_-$ be the subbundle of $\text{End}(E)$ corresponding to the negative eigenspaces of $i\beta$ and let $\rho_x^- : \Omega^0(\text{End}(E)_-) \rightarrow \Omega^{0,1}(\text{End}(E)_-) \oplus \Omega^{1,0}(\text{End}(E)_-)$ be the restriction of the infinitesimal action to the negative eigenspaces. Then

$$\text{im } \rho_x^- = \text{im } \rho_x \cap \Omega^{0,1}(\text{End}(E)_-) \oplus \Omega^{1,0}(\text{End}(E)_-)$$

and

$$(2.9) \quad \ker(\rho_x^-)^* \supseteq \ker \rho_x^* \cap \Omega^{0,1}(\text{End}(E)_-) \oplus \Omega^{1,0}(\text{End}(E)_-)$$

Since $\text{im } \rho_x \oplus \ker \rho_x^* \cong \Omega^{0,1}(\text{End}(E)) \oplus \Omega^{1,0}(\text{End}(E))$ by [51, Lem. 4.9] then

$$\begin{aligned} \Omega^{0,1}(\text{End}(E)_-) \oplus \Omega^{1,0}(\text{End}(E)_-) &= (\text{im } \rho_x \oplus \ker \rho_x^*) \cap (\Omega^{0,1}(\text{End}(E)_-) \oplus \Omega^{1,0}(\text{End}(E)_-)) \\ &\subseteq (\text{im } \rho_x^- \oplus \ker(\rho_x^-)^*) \subseteq \Omega^{0,1}(\text{End}(E)_-) \oplus \Omega^{1,0}(\text{End}(E)_-) \end{aligned}$$

and so (2.9) must be an equality, therefore $\Omega^{0,1}(\text{End}(E)_-) \oplus \Omega^{1,0}(\text{End}(E)_-) \cong \text{im } \rho_x^- \oplus \ker(\rho_x^-)^*$. Therefore the function

$$\begin{aligned} \psi^- : (\ker \rho_x^-)^\perp \times \ker(\rho_x^-)^* &\rightarrow \Omega^{0,1}(\text{End}(E)_-) \oplus \Omega^{1,0}(\text{End}(E)_-) \\ (u, \delta x) &\mapsto e^u \cdot (x + \delta x) \end{aligned}$$

is a local diffeomorphism at 0. If $\delta x \in S_x^-$ then $x + \delta x \in \mathcal{B}$, and so $e^u \cdot (x + \delta x) \in \mathcal{B}$, since the complex gauge group preserves the space of Higgs bundles. Conversely, if $e^u \cdot (x + \delta x) \in \mathcal{B}$ then $x + \delta x \in \mathcal{B}$ and so $\delta x \in S_x^-$. Therefore ψ restricts to a local homeomorphism $(\ker \rho_x^-)^\perp \times S_x^- \rightarrow \mathcal{B} \cap (\Omega^{0,1}(\text{End}(E)_-) \oplus \Omega^{1,0}(\text{End}(E)_-))$. \square

The next two results concern a sequence of points $g_t \cdot z$ in a $\mathcal{G}^{\mathbb{C}}$ orbit which approach a critical point x in the L_k^2 norm and for which $\text{YMH}(z) < \text{YMH}(x)$. Since x is critical and $\text{YMH}(z) < \text{YMH}(x)$ then $x \in \overline{\mathcal{G}^{\mathbb{C}} \cdot z} \setminus \mathcal{G}^{\mathbb{C}} \cdot z$, and therefore $\|g_t\|_{L_{k+1}^2} \rightarrow \infty$. The result below shows that the C^0 norm of the function $\sigma(h_t) = \text{Tr}(h_t) + \text{Tr}(h_t^{-1}) - 2 \text{rank}(E)$ must also blow up.

Lemma 2.11. *Let $x \in \mathcal{B}$ be a critical point of YMH and let $z \in \mathcal{B}$ be any point such that $\text{YMH}(z) < \text{YMH}(x)$. Suppose that there exists a sequence of gauge transformations $g_t \in \mathcal{G}^{\mathbb{C}}$ such that $g_t \cdot z \rightarrow x$ in L_k^2 . Then the change of metric $h_t = g_t^* g_t$ satisfies $\sup_X \sigma(h_t) \rightarrow \infty$.*

Proof. Let U be the neighbourhood of x from Lemma 2.8. Since $g_t \cdot z \rightarrow x$, then there exists T such that $g_t \cdot z \in U$ for all $t \geq T$. Therefore there exists f_t in a neighbourhood of the identity in $\mathcal{G}^{\mathbb{C}}$ such that $f_t \cdot g_t \cdot z \in S_x$. The uniqueness of the decomposition from the slice theorem shows that if $t > T$, then $f_t \cdot g_t \cdot z = f_{t,T} \cdot f_T \cdot g_T \cdot z$ with $f_{t,T} \in \mathcal{G}_x^{\mathbb{C}}$. Therefore $t \rightarrow \infty$ implies that $f_{t,T}$ diverges in $\mathcal{G}_x^{\mathbb{C}}$. Fix a point p on the surface X , and let $\mathcal{G}_0^{\mathbb{C}}$ be the based gauge group consisting of complex gauge transformations that are the identity at p . Recall that $\mathcal{G}_0^{\mathbb{C}}$ is a normal subgroup of $\mathcal{G}^{\mathbb{C}}$ and we have the following short exact sequence of groups (cf. [1, Sec. 13])

$$1 \rightarrow \mathcal{G}_0^{\mathbb{C}} \rightarrow \mathcal{G}^{\mathbb{C}} \rightarrow \text{GL}(n, \mathbb{C}) \rightarrow 1.$$

Since $\mathcal{G}_0^{\mathbb{C}}$ acts freely on the space of connections (and hence on \mathcal{B}), then restriction to the fibre over p defines a bijective correspondence between $\mathcal{G}_x^{\mathbb{C}} \subset \mathcal{G}^{\mathbb{C}}$ and a subgroup of $\text{GL}(n, \mathbb{C})$ via the exact sequence above. Therefore $f_{t,T}$ diverges in $\mathcal{G}_x^{\mathbb{C}}$ implies that the restriction of $f_{t,T}$ to the fibre over p diverges in $\text{GL}(n, \mathbb{C})$, and so the C^0 norm of $f_{t,T}$ diverges to ∞ , and hence the same is true for $g_t = f_t^{-1} \cdot f_{t,T} \cdot f_T \cdot g_T \cdot z$ since g_T is fixed and both f_t and f_T are contained in a fixed neighbourhood of the identity in $\mathcal{G}^{\mathbb{C}}$. Therefore $\sup_X \sigma(h_t) \rightarrow \infty$. \square

Corollary 2.12. *Let x be a critical point of YMH. Then for each neighbourhood V of x in the L_k^2 topology on \mathcal{B} and each constant $C > 0$, there exists a neighbourhood U of x such that if $z \notin V$ and $\text{YMH}(z) < \text{YMH}(x)$, then $y = g \cdot z$ with $h = g^* g$ satisfying $\sup_X \sigma(h) \leq C$ implies that $y \notin U$.*

Proof. If no such neighbourhood U exists, then we can construct a sequence $y_t = g_t \cdot z$ converging to x in L_k^2 such that $h_t = g_t^* g_t$ satisfies $\sup_X \sigma(h_t) \leq C$ for all t , however this contradicts the previous lemma. \square

2.3. Modifying the YMH flow in a neighbourhood of a critical point. Let x be a critical point, let $\beta = \mu(x) = *(F_A + [\phi, \phi^*])$, and let $\mathcal{G}_*^{\mathbb{C}}$ be the subgroup defined in the previous section. In this section we explain how to modify the YMH flow near x so that the gauge transformation generating the flow is contained in $\mathcal{G}_*^{\mathbb{C}}$. The reason for modifying the flow is so that we can apply the distance-decreasing formula of Lemma 2.19, which is used for the convergence result of Section 3.2.

Let U be a \mathcal{G} -invariant neighbourhood of x such that U is homeomorphic to a neighbourhood of $[\text{id}, 0, 0]$ in $\mathcal{G} \times_{\mathcal{G}_\beta} ((\ker \rho_x)_*^\perp \times S_x)$ by Proposition 2.9. Let $V \subset U$ be the image of $(\ker \rho_x)_*^\perp \times S_x$ under the homeomorphism from Proposition 2.9. For each $y \in V$, let $\gamma_-(y)$ be the component of $i\mu(y)$ in $\Omega^0(\text{End}(E)_-)$. Since μ is \mathcal{G} -equivariant then we can extend γ_- equivariantly from V to all of U using the action of \mathcal{G} . Define the map $\gamma : U \rightarrow \text{Lie}(\mathcal{G})$ by

$$(2.10) \quad \gamma(y) = \gamma_-(y) - \gamma_-(y)^*$$

so that $-i\mu(y) + \gamma(y) \in \text{Lie}(\mathcal{G}_*^{\mathbb{C}})$.

Definition 2.13. The *modified flow* with initial condition $y_0 \in U$ is the solution to

$$(2.11) \quad \frac{dy}{dt} = -I\rho_x(\mu(y)) + \rho_x(\gamma(y)).$$

More explicitly, on the space of Higgs bundles $y = (\bar{\partial}_A, \phi)$ satisfies

$$\begin{aligned} \frac{\partial A}{\partial t} &= i\bar{\partial}_A * (F_A + [\phi, \phi^*]) - \bar{\partial}_A \gamma(\bar{\partial}_A, \phi) \\ \frac{\partial \phi}{\partial t} &= i[\phi, *(F_A + [\phi, \phi^*])] - [\phi, \gamma(\bar{\partial}_A, \phi)] \end{aligned}$$

In analogy with (2.3), the modified flow is generated by the action of the gauge group $y_t = g_t \cdot y_0$, where g_t satisfies the equation

$$(2.12) \quad \frac{\partial g_t}{\partial t} g_t^{-1} = -i\mu(g_t \cdot y_0) + \gamma(g_t \cdot y_0), \quad g_0 = \text{id}.$$

As before, let $V \subset U$ be the image of $(\ker \rho_x)_*^\perp \times S_x$ under the homeomorphism from the slice theorem (Proposition 2.9). Note that if $y_0 \in V$ then $\frac{\partial g_t}{\partial t} g_t^{-1} \in \text{Lie}(\mathcal{G}_*^{\mathbb{C}})$, so $g_t \in \mathcal{G}_*^{\mathbb{C}}$ and the solution to the modified flow remains in V for as long as it remains in the neighbourhood U .

Lemma 2.14. *Let $y_t = g_t \cdot y_0$ be the solution to the YMH flow (2.3) with initial condition y_0 . Then there exists $s_t \in \mathcal{G}$ solving the equation*

$$(2.13) \quad \frac{ds}{dt} s_t^{-1} = \gamma(s_t \cdot y_t), \quad s_0 = \text{id}$$

such that $\tilde{y}_t = s_t \cdot y_t$ is a solution to the modified flow equation (2.11) with initial condition y_0 .

Proof. Since γ is \mathcal{G} -equivariant then (2.13) reduces to

$$\frac{ds}{dt} s_t^{-1} = \text{Ad}_{s_t} \gamma(y_t).$$

Since $\gamma(y_t) \in \text{Lie}(\mathcal{G})$ is already defined by the gradient flow y_t , then this equation reduces to solving an ODE on the fibres of the bundle, and therefore existence of solutions follows from ODE existence theory.

Let $\tilde{g}_t = s_t \cdot g_t$. A calculation shows that

$$\begin{aligned} \frac{d\tilde{g}_t}{dt} \tilde{g}_t^{-1} &= \frac{ds}{dt} s_t^{-1} + \text{Ad}_{s_t} \left(\frac{dg}{dt} g_t^{-1} \right) \\ &= \gamma(s_t \cdot y_t) - i \text{Ad}_{s_t} \mu(y_t) \\ &= \gamma(\tilde{y}_t) - i\mu(\tilde{y}_t) \\ &= \gamma(\tilde{g}_t \cdot y_0) - i\mu(\tilde{g}_t \cdot y_0), \end{aligned}$$

and so $\tilde{y}_t = \tilde{g}_t \cdot y_0 = s_t \cdot y_t$ is a solution to the modified flow (2.11) with initial condition y_0 . \square

As a corollary, we see that the change of metric is the same for the YMH flow (2.3) and the modified flow (2.12).

Corollary 2.15. *Let $y_t = g_t \cdot y_0$ be a solution to the Yang-Mills-Higgs flow equation (2.3) and $\tilde{y}_t = \tilde{g}_t \cdot y_0$ be a solution to the modified flow equation (2.12). Then $h_t = g_t^* g_t = \tilde{g}_t^* \tilde{g}_t$.*

Finally, we prove that convergence for the upwards YMH flow implies convergence for the modified flow.

Lemma 2.16. *Let x be a critical point and let $y_0 \in W_x^-$. Then the modified flow with initial condition y_0 exists for all $t \in (-\infty, 0]$ and converges in the C^∞ topology to a point in $\mathcal{G} \cdot x$.*

Proof. Let y_t be the YMH flow with initial condition y_0 and $\tilde{y}_t = s_t \cdot y_t$ the modified flow. By the definition of W_x^- the YMH flow exists for all $t \in (-\infty, 0]$ and $y_t \rightarrow x$ in the C^∞ topology. Existence of the modified flow then follows from Lemma 2.14. Proposition 2.21 shows that $y_t \rightarrow x$ exponentially in L_k^2 for all k , and so the same is true for $\gamma(y_t)$. Therefore the length of the modified flow line satisfies

$$\begin{aligned} \int_{-\infty}^0 \|I\rho_{\tilde{y}_t}(\mu(\tilde{y}_t)) - \rho_{\tilde{y}_t}(\gamma(\tilde{y}_t))\|_{L_k^2} dt &= \int_{-\infty}^0 \|I\rho_{y_t}(\mu(y_t)) - \rho_{y_t}(\gamma(y_t))\|_{L_k^2} dt \\ &\leq \int_{-\infty}^0 \|\rho_{y_t}(\mu(y_t))\|_{L_k^2} dt + \int_{-\infty}^0 \|\rho_{y_t}(\gamma(y_t))\|_{L_k^2} dt \end{aligned}$$

which is finite since the length $\int_{-\infty}^0 \|\rho_{y_t}(\mu(y_t))\|_{L_k^2} dt$ of the YMH flow line is finite, y_t is bounded and $\gamma(y_t) \rightarrow 0$ exponentially. This is true for all k , and so the modified flow converges in the C^∞ topology. \square

2.4. Preliminary estimates for the YMH flow in a neighbourhood of a critical point. Given eigenvalues for the adjoint action of $i\beta$ on $\text{Lie}(\mathcal{G}^{\mathbb{C}})$ labelled by $\lambda_1 \leq \dots \leq \lambda_k < 0 \leq \lambda_{k+1} \leq \dots$ (with respect to (2.6), the negative eigenvalues correspond to $\lambda_\ell = \nu_i - \nu_j$ for $i < j$), then for any $y \in S_x^-$ and any norm, we have the Lipschitz bounds

$$(2.14) \quad e^{\lambda_1 t} \|y - x\| \leq \|e^{i\beta t} \cdot y - x\| \leq e^{\lambda_k t} \|y - x\|.$$

Lemma 2.17. *For any critical point x there exists $C > 0$ such that for any $y \in S_x^-$, we have $\|\mu(y) - \beta\|_{C^0} \leq C\|y - x\|_{C^0}^2$.*

Proof. Let $y \in S_x^-$ and define $\delta y := y - x \in V \cong T_x V$. Then the defining equation for the moment map shows that for all $v \in \mathfrak{k}$, we have

$$d\mu_x(\delta y) \cdot v = \omega(\rho_x(v), \delta y) = \langle I\rho_x(v), \delta y \rangle$$

By the definition of the slice, each $\delta y \in S_x^-$ is orthogonal to the infinitesimal action of $\mathcal{G}^{\mathbb{C}}$ at x , and so $\langle I\rho_x(v), \delta y \rangle = 0$ for all $v \in \mathfrak{k}$. Therefore $d\mu_x(\delta y) = 0$. Since the moment map $\mu(\bar{\partial}_A, \phi) = F_A + [\phi, \phi^*]$ is quadratic, then we have

$$\|\mu(y) - \mu(x)\|_{C^0} \leq \|d\mu_x(\delta y)\|_{C^0} + C\|\delta y\|_{C^0}^2 = C\|\delta y\|_{C^0}^2.$$

Since the moment map is \mathcal{G} -equivariant and the norms above are all \mathcal{G} -invariant, then the constant C is independent of the choice of critical point in the orbit $\mathcal{G} \cdot x$. \square

Given $g \in \mathcal{G}^{\mathbb{C}}$, let g^* denote the adjoint with respect to the Hermitian metric on E and let \mathcal{G} act on $\mathcal{G}^{\mathbb{C}}$ by left multiplication. In every equivalence class of the space of metrics $\mathcal{G}^{\mathbb{C}}/\mathcal{G}$ there is a unique positive definite self-adjoint section h , which we use from now on to represent elements of $\mathcal{G}^{\mathbb{C}}/\mathcal{G}$. Given $h = g^*g \in \mathcal{G}^{\mathbb{C}}/\mathcal{G}$, define $\mu_h : \mathcal{B} \rightarrow \Omega^0(\text{End}(E)) \cong \text{Lie}(\mathcal{G}^{\mathbb{C}})$ by

$$(2.15) \quad \mu_h(y) = \text{Ad}_{g^{-1}}(\mu(g \cdot y)).$$

Since the moment map is \mathcal{G} -equivariant, then for any $k \in \mathcal{G}$ we have

$$\text{Ad}_{g^{-1}} \text{Ad}_{k^{-1}} (\mu(k \cdot g \cdot y)) = \text{Ad}_{g^{-1}} (\mu(g \cdot y))$$

and so μ_h is well-defined on $\mathcal{G}^{\mathbb{C}}/\mathcal{G}$. The length of a geodesic in the space of positive definite Hermitian matrices is computed in [27, Ch. VI.1]. Following [8, Prop. 13] (see also [45, Prop. 6.3]), it is more convenient to define the distance function $\sigma : \mathcal{G}^{\mathbb{C}}/\mathcal{G} \rightarrow \mathbb{R}$

$$(2.16) \quad \sigma(h) = \text{Tr } h + \text{Tr } h^{-1} - 2 \text{rank}(E).$$

As explained in [8], the function $\sup_X \sigma$ is not a norm in the complete metric space $\mathcal{G}^{\mathbb{C}}/\mathcal{G}$, however we do have $h_t \xrightarrow{C^0} h_\infty$ in $\mathcal{G}^{\mathbb{C}}/\mathcal{G}$ if and only if $\sup_X \sigma(h_t h_\infty^{-1}) \rightarrow 0$. Note that if $h_1 = g_1^* g_1$ and $h_2 = g_2^* g_2$, then

$$(2.17) \quad \sigma(h_1 h_2^{-1}) = \sigma(g_1^* g_1 g_2^{-1} (g_2^*)^{-1}) = \sigma((g_1 g_2^{-1})^* g_1 g_2^{-1}).$$

Recall from [8], [45] that we have the following distance-decreasing formula for a solution to the downwards YMH flow. Since the change of metric is the same for the modified flow by Corollary 2.15, then (2.18) is also valid for the modified flow.

Lemma 2.18. *Let $y_1, y_2 \in \mathcal{B}$ and suppose that $y_1 = g_0 \cdot y_2$ for $g_0 \in \mathcal{G}^{\mathbb{C}}$. For $j = 1, 2$, define $y_j(t)$ to be the solution of the YMH flow (2.4) with initial condition y_j . Define g_t by $y_1(t) = g_t \cdot y_2(t)$ and let $h_t = g_t^* g_t$ be the associated change of metric. Then*

$$(2.18) \quad \left(\frac{\partial}{\partial t} + \Delta \right) \sigma(h_t) \leq 0.$$

Since $\text{Lie}(\mathcal{G}_*^{\mathbb{C}}) = \Omega^0(\text{End}(E))_0 \oplus \Omega^0(\text{End}(E))_+$ and the adjoint action of $e^{-i\beta t}$ is the identity on $\Omega^0(\text{End}(E))_0$ and strictly contracting on $\Omega^0(\text{End}(E))_+$, then we have the following lemma which is used in Section 3.2.

Lemma 2.19. *Given any $g_0 \in \mathcal{G}_*^{\mathbb{C}}$, let $g_t = e^{-i\beta t} g_0 e^{i\beta t}$ and $h_t = g_t^* g_t$. Then $\frac{\partial}{\partial t} \sigma(h_t) \leq 0$.*

In the proof of the distance-decreasing formula for the Yang-Mills flow [8, Prop. 13], Donaldson proves that for any holomorphic structure $\bar{\partial}_A$ and any complex gauge transformation g , the change of metric $h = g^* g$ satisfies

$$-2i\Lambda \text{Tr}(h(g^{-1}(F_{g \cdot A})g - F_A)) = -2i\Lambda \text{Tr}(\bar{\partial}_A \partial_A h - (\bar{\partial}_A h)h(\partial_A h)) \leq -\Delta \text{Tr}(h).$$

Note that the notation here differs slightly from that in [8, Prop. 13] which considers a fixed holomorphic structure and a varying Hermitian metric, while here we consider a fixed Hermitian metric and a varying holomorphic structure. As explained on [8, p5] the two viewpoints are equivalent. A similar calculation shows that

$$2i\Lambda \text{Tr}(h^{-1}(g^{-1}(F_{g \cdot A})g - F_A)) \leq -\Delta \text{Tr}(h^{-1}).$$

Analogous calculations for a Higgs pair $y = (\bar{\partial}_A, \phi)$ using $\mu(y) = \Lambda(F_A + [\phi, \phi^*])$ instead of the curvature ΛF_A gives us the following result for the Yang-Mills-Higgs flow, which is used in the proof of Lemma 3.2.

Lemma 2.20.

$$\begin{aligned} -2i \operatorname{Tr}((\mu_h(y) - \mu(y))h) + \Delta \operatorname{Tr}(h) &\leq 0 \\ 2i \operatorname{Tr}((\mu_h(y) - \mu(y))h^{-1}) + \Delta \operatorname{Tr}(h^{-1}) &\leq 0. \end{aligned}$$

2.5. Exponential convergence of the backwards flow. In this section we prove that if a solution to the backwards YMH flow converges to a critical point, then it must do so exponentially in each Sobolev norm.

Proposition 2.21. *Let y_t be a solution to the YMH flow (2.4) such that $\lim_{t \rightarrow -\infty} y_t = x$. Then for each positive integer k there exist positive constants C_1 and η such that $\|y_t - x\|_{L_k^2} \leq C_1 e^{\eta t}$ for all $t \leq 0$.*

The proof of the proposition reduces to the following lemmas. First recall from the slice theorem that there is a unique decomposition

$$y = e^u \cdot (x + z)$$

for $u \in (\ker \rho_x)^\perp$ and $z \in S_x$. We can further decompose $z = z_{\geq 0} + z_-$, where $z_- \in S_x^-$ is the component of z in the negative slice and $z_{\geq 0} = z - z_-$. At the critical point x we have the decomposition $\operatorname{End}(E) \cong \operatorname{End}(E)_+ \oplus \operatorname{End}(E)_0 \oplus \operatorname{End}(E)_-$ according to the eigenspaces of $i\beta$ (cf. Sec. 2.2). Then with respect to this decomposition $z_{\geq 0}$ is the component of z in $\Omega^{0,1}(\operatorname{End}(E)_+) \oplus \operatorname{End}(E)_0 \oplus \Omega^{1,0}(\operatorname{End}(E)_+ \oplus \operatorname{End}(E)_0)$ and z_- is the component in $\Omega^{0,1}(\operatorname{End}(E)_-) \oplus \Omega^{1,0}(\operatorname{End}(E)_-)$. In terms of the action of $\beta = \mu(x)$ we have $\lim_{t \rightarrow \infty} e^{i\beta t} \cdot z_- = 0$ and $\lim_{t \rightarrow \infty} e^{-i\beta t} \cdot z_{\geq 0} = z_0$, where z_0 is the component of z in $\Omega^{0,1}(\operatorname{End}(E)_0) \oplus \Omega^{1,0}(\operatorname{End}(E)_0)$. Note that if $y = e^u \cdot (x + z)$ is a Higgs bundle, then $x + z$ is a Higgs bundle since $e^u \in \mathcal{G}^\mathbb{C}$ preserves the space of Higgs bundles, however $x + z_{\geq 0}$ may not be a Higgs bundle as the pair $(\bar{\partial}_{A_{\geq 0}}, \phi_{\geq 0})$ representing $x + z_{\geq 0}$ may not satisfy $\bar{\partial}_{A_{\geq 0}} \phi_{\geq 0} = 0$. Even though $\phi_{\geq 0}$ may not be holomorphic, we can still apply the principle that curvature decreases in subbundles and increases in quotient bundles and follow the same idea as [1, Sec. 8 & 10] to prove the following lemma.

Lemma 2.22. (1) $\operatorname{grad} \operatorname{YMH}(e^u \cdot (x + z_{\geq 0}))$ is tangent to the set $\{z_- = 0\}$.
 (2) $\operatorname{YMH}(e^u \cdot (x + z_{\geq 0})) \geq \operatorname{YMH}(x)$.

Proof. By definition of z_- as the component of $z \in S_x$ in $\Omega^{0,1}(\operatorname{End}(E)_-) \oplus \Omega^{1,0}(\operatorname{End}(E)_-)$, the subset $\{z_- = 0\}$ consists of all pairs $(\bar{\partial}_A, \phi)$ such that (up to complex gauge transformations) the bundle E admits a filtration $E_1 \subset \dots \subset E_n = E$ by holomorphic subbundles of the same Harder-Narasimhan type as x which are also preserved by ϕ . Note that this does not require ϕ to be holomorphic since we have constructed the filtration explicitly using the decomposition $\operatorname{End}(E) \cong \operatorname{End}(E)_+ \oplus \operatorname{End}(E)_0 \oplus \operatorname{End}(E)_-$. Since $\operatorname{grad} \operatorname{YMH}(e^u \cdot (x + z_{\geq 0}))$ is tangent to the complex gauge orbit through $e^u \cdot (x + z_{\geq 0})$, and the existence of a filtration of a given type is preserved by the action of the complex gauge group, then $\operatorname{grad} \operatorname{YMH}(e^u \cdot (x + z_{\geq 0}))$ is tangent to the set $\{z_- = 0\}$.

Below are the details of the proof of the second part of the lemma for the case of a two-step filtration. This can be generalised to an arbitrary filtration in the same way as [1, Sec. 8].

Let ξ be a convex invariant function on the Lie algebra of the structure group. For the following, we only need to consider the special case of $\xi(X) = \operatorname{Tr}(XX^*)$ but it is worth noting that the entire argument works for any convex invariant ξ . Consider a critical point x such that the C^∞ bundle E decomposes as a direct

sum $E_1 \oplus E_2$ and the Higgs bundle x is the direct sum of Hermitian-Yang-Mills-Higgs pairs on E_1 and E_2 . Then $\text{YMH}(x) = \int_X \xi(\mu) \text{dvol}$, where

$$\mu = \begin{pmatrix} \frac{\deg E_1}{\text{rank } E_1} \cdot \text{id} & 0 \\ 0 & \frac{\deg E_2}{\text{rank } E_2} \cdot \text{id} \end{pmatrix}$$

Consider an extension

$$0 \rightarrow (\bar{\partial}_{A_1}, \phi_1) \rightarrow (\bar{\partial}_A, \phi) \rightarrow (\bar{\partial}_{A_2}, \phi_2) \rightarrow 0$$

for which $(\bar{\partial}_{A_1}, \phi_1)$ and $(\bar{\partial}_{A_2}, \phi_2)$ are arbitrary Higgs pairs on E_1 and E_2 . Locally, we can write the holomorphic structure $\bar{\partial}_A$ and the Higgs field ϕ as follows

$$\bar{\partial}_A = \begin{pmatrix} \bar{\partial}_{A_1} & a \\ 0 & \bar{\partial}_{A_2} \end{pmatrix}, \quad \partial_{A,h} = \begin{pmatrix} \partial_{A_1,h} & 0 \\ -a^* & \partial_{A_2,h} \end{pmatrix}, \quad \phi = \begin{pmatrix} \phi_1 & \varphi \\ 0 & \phi_2 \end{pmatrix}, \quad \phi^{*h} = \begin{pmatrix} \phi_1^* & 0 \\ \varphi^* & \phi_2^* \end{pmatrix}$$

Note that we are not assuming that $\bar{\partial}_A \phi = 0$. From now on we suppress the notation for the metric h . We then have (cf. [1, (8.14)])

$$F_A + [\phi, \phi^*] = \begin{pmatrix} F_{A_1} - a \wedge a^* & d_A a \\ -d_A a^* & F_{A_2} - a^* \wedge a \end{pmatrix} + \begin{pmatrix} [\phi_1, \phi_1^*] + \varphi \wedge \varphi^* & \varphi \wedge \phi_2^* + \phi_1^* \wedge \varphi \\ \phi_2 \wedge \varphi^* + \varphi^* \wedge \phi_1 & [\phi_2, \phi_2^*] + \varphi^* \wedge \varphi \end{pmatrix}$$

Now for $j = 1, 2$, let f_j, α_j be scalar $\text{rank}(E_j) \times \text{rank}(E_j)$ matrix valued functions (cf. [1, (8.15)]) such that

$$\text{Tr } f_j = * \text{Tr}(F_{A_j} + [\phi_j, \phi_j^*]), \quad \text{Tr } \alpha_1 = * \text{Tr}(a \wedge a^* - \varphi \wedge \varphi^*) = - * \text{Tr}(a^* \wedge a - \varphi^* \wedge \varphi) = - \text{Tr } \alpha_2$$

Then for any convex invariant function ξ , the results of [1, Sec. 12] show that

$$\xi(* (F_A + [\phi, \phi^*])) \geq \xi \begin{pmatrix} f_1 - \alpha_1 & 0 \\ 0 & f_2 - \alpha_2 \end{pmatrix}$$

If we normalise so that $\text{vol}(X) = 1$ then combining the above inequality with Jensen's inequality gives us

$$\text{YMH}(\bar{\partial}_A, \phi) = \int_X \xi(* (F_A + [\phi, \phi^*])) \text{dvol} \geq \xi \left(\int_X \begin{pmatrix} f_1 - \alpha_1 & 0 \\ 0 & f_2 - \alpha_2 \end{pmatrix} \text{dvol} \right)$$

Since the Lie bracket $[\phi, \phi^*]$ is traceless, then the degree of E_j is

$$k_j = \frac{i}{2\pi} \int_X \text{Tr}(F_{A_j} + [\phi_j, \phi_j^*]) = \frac{i}{2\pi} \int_X \text{Tr}(f_j) \text{dvol}$$

Now we note that $-i * \text{Tr}(a \wedge a^*) \geq 0$ and $i * \text{Tr}(\varphi \wedge \varphi^*) \geq 0$. The difference in sign is because $a \in \Omega^{0,1}$ and $\varphi \in \Omega^{1,0}$. For completeness, we give all the details. Choose a local coordinate $z = x + iy$ such that $dx \wedge dy = \text{dvol}$. First note that $d\bar{z} \wedge dz = 2i dx \wedge dy = 2i \text{dvol}$. Therefore $-i * (d\bar{z} \wedge dz) = 2$ and $i * (dz \wedge d\bar{z}) = 2$. Writing $a = g_1 d\bar{z}$ in a local neighbourhood, we have

$$-i * \text{Tr}(a \wedge a^*) = -i * \text{Tr}(g_1 g_1^* d\bar{z} \wedge dz) = 2 \text{Tr}(g_1 g_1^*) \geq 0$$

and writing $\varphi = g_2 dz$ in a local neighbourhood we have

$$i * \text{Tr}(\varphi \wedge \varphi^*) = i * \text{Tr}(g_2 g_2^* dz \wedge d\bar{z}) = 2 \text{Tr}(g_2 g_2^*) \geq 0$$

For $j = 1, 2$, define $b_j = -\frac{i}{2\pi} \int_X \alpha_j$. Then the above calculations show that b_1 is a scalar matrix with non-negative entries, and b_2 is a scalar matrix with non-positive entries.

Putting all of this together, we have

$$\int_X \begin{pmatrix} f_1 - \alpha_1 & 0 \\ 0 & f_2 - \alpha_2 \end{pmatrix} \text{dvol} = -2\pi i(\mu + b)$$

where

$$\mu = \begin{pmatrix} \frac{\deg E_1}{\text{rank } E_1} \cdot \text{id} & 0 \\ 0 & \frac{\deg E_2}{\text{rank } E_2} \cdot \text{id} \end{pmatrix}, \quad b = \begin{pmatrix} b_1 & 0 \\ 0 & b_2 \end{pmatrix}$$

Since $\text{Tr } \alpha_1 = -\text{Tr } \alpha_2$, then $\text{Tr } b = 0$ and so the inequalities in [1, Sec. 12] for convex invariant functions show that $\|F_A + [\phi, \phi^*]\|_{L_2}^2 \geq \xi(\mu) = \text{YMH}(x)$. \square

The next lemma shows that the component in the negative slice is decreasing exponentially.

Lemma 2.23. *Let $y_t = e^u \cdot (x + z_{\geq 0} + z_-)$ be a solution to the YMH flow such that $\lim_{t \rightarrow -\infty} y_t = x$. Then there exist positive constants K_1 and K_2 such that $\|z_-\|_{L_2}^2 \leq K_1 e^{K_2 t}$ for all $t \leq 0$.*

Proof. The proof follows the idea of [26, Sec. 10]. The downwards gradient flow equation for z_- is

$$\frac{\partial z_-}{\partial t} = Lz_- + N_-(u, z_{\geq 0}, z_-)$$

where L is a linear operator and the derivative of N_- vanishes at the origin. Since z_- is orthogonal to the $\mathcal{G}^{\mathbb{C}}$ orbit through x , then the following calculation shows that the linear part satisfies $e^{Lt} z_- = e^{-i\beta t} \cdot z_-$.

Let $(\bar{\partial}_{A_\infty}, \phi_\infty)$ be the critical point x and $(\bar{\partial}_A, \phi) = (\bar{\partial}_{A_\infty}, \phi_\infty) + (a, \varphi)$ be a point in the slice S_x . The condition that (a, φ) is orthogonal to the $\mathcal{G}^{\mathbb{C}}$ orbit through $(\bar{\partial}_A, \phi)$ is given by

$$\bar{\partial}_{A_\infty}^* a - \bar{*}[\phi, \bar{*}\varphi] = 0$$

We can rewrite this orthogonality condition as

$$i * \partial_A a + i * [\phi^*, \varphi] = 0 \quad \Leftrightarrow \quad i * \bar{\partial}_A a^* - i * [\phi, \varphi^*] = 0$$

Subtracting the first equation from the second and using the fact that the Hodge star is an isomorphism $\Omega^2 \cong \Omega^0$ gives us

$$d_A(a - a^*) + [\phi, \varphi^*] + [\varphi, \phi^*] = 0.$$

Note also that the curvature can be written as

$$F_{A+a} + [\phi + \varphi, (\phi + \varphi)^*] = F_A + [\phi, \phi^*] + d_A(a - a^*) + [\varphi, \phi^*] + [\phi, \varphi^*] + (\text{higher order terms})$$

Therefore the slice equations imply that the linearisation of the curvature vanishes in the direction of a . Substituting this into the gradient flow equations and using the fact that $(\bar{\partial}_A, \phi)$ is a critical point gives us

$$\begin{aligned} \frac{\partial a}{\partial t} &= i[a, *(F_A + [\phi, \phi^*])] + i\bar{\partial}_A * (d_A(a - a^*) + [\phi, \varphi^*] + [\varphi, \phi^*]) + (\text{higher order terms}) \\ \frac{\partial \varphi}{\partial t} &= i[\varphi, *(F_A + [\phi, \phi^*])] + i[\phi, (d_A(a - a^*) + [\phi, \varphi^*] + [\varphi, \phi^*])] + (\text{higher order terms}) \end{aligned}$$

Therefore at a point in the slice, the linear component of the gradient flow equations is given by $L(a, \varphi) = (-i[\beta, a], -i[\beta, \varphi])$, where $\beta = *(F_A + [\phi, \phi^*])$. Note that this is the tangent vector to the curve $e^{-i\beta t} \cdot (a, \varphi)$ at $t = 0$, and therefore if (a, φ) is in the negative slice then the action of $-i\beta$ has strictly positive eigenvalues (i.e. the downwards flow increases the component in the negative slice).

Since z_- is in the negative slice then there exists $\lambda_{min} > 0$ such that $\langle Lz_-, z_- \rangle_{L_1^2} \geq \lambda_{min} \|z_-\|_{L_1^2}^2$. Now Lemma 2.22 shows that the YMH flow preserves the set $\{z_- = 0\}$, and so $N_-(u, z_{\geq 0}, 0) = 0$. Since N_- is C^1 with vanishing derivative at the origin then for all $\varepsilon > 0$ there exists $\delta > 0$ such that if $\|y_t - x\|_{L_1^2} < \delta$ then

$$\|N_-(u, z_{\geq 0}, z_-)\|_{L_1^2} \leq \varepsilon \|z_-\|_{L_1^2}$$

Therefore

$$\frac{1}{2} \frac{\partial}{\partial t} \|z_-\|_{L_1^2}^2 = \langle Lz_-, z_- \rangle_{L_1^2} + \langle N_-(u, z_{\geq 0}, z_-), z_- \rangle_{L_1^2} \geq (\lambda_{min} - \varepsilon) \|z_-\|_{L_1^2}^2,$$

and so if $\varepsilon > 0$ is small enough (e.g. $\varepsilon < \frac{1}{2}\lambda_{min}$) then there exist positive constants K_1 and K_2 such that $\|z_-\|_{L_1^2}^2 \leq K_1 e^{K_2 t}$ for all $t \leq 0$. \square

The next lemma shows that the difference $\text{YMH}(x) - \text{YMH}(y_t)$ is decreasing exponentially.

Lemma 2.24. *Let $y_t = e^u \cdot (x + z_{\geq 0} + z_-)$ be a solution to the YMH flow such that $\lim_{t \rightarrow -\infty} y_t = x$. Then there exist positive constants K'_1 and K'_2 such that*

$$\text{YMH}(x) - \text{YMH}(e^u \cdot (x + z_{\geq 0} + z_-)) \leq K'_1 e^{K'_2 t}$$

for all $t \leq 0$.

Proof. Recall that the Morse-Kirwan condition from Lemma 2.22 implies

$$\text{YMH}(e^u \cdot (x + z_{\geq 0})) - \text{YMH}(x) \geq 0$$

Since x is a critical point of YMH, then for all $\varepsilon > 0$ there exists $\delta > 0$ such that if $\|y_t - x\|_{L_1^2} < \delta$ we have

$$\text{YMH}(e^u \cdot (x + z_{\geq 0} + z_-)) - \text{YMH}(e^u \cdot (x + z_{\geq 0})) \geq -\varepsilon \|z_-\|_{L_1^2}.$$

Therefore

$$\begin{aligned} \text{YMH}(e^u \cdot (x + z_{\geq 0} + z_-)) - \text{YMH}(x) &= \text{YMH}(e^u \cdot (x + z_{\geq 0} + z_-)) - \text{YMH}(e^u \cdot (x + z_{\geq 0})) \\ &\quad + \text{YMH}(e^u \cdot (x + z_{\geq 0})) - \text{YMH}(x) \\ &\geq -\varepsilon \|z_-\|_{L_1^2} \geq -\varepsilon \sqrt{K_1} e^{\frac{1}{2} K_2 t} \end{aligned}$$

Since $\text{YMH}(e^u \cdot (x + z_{\geq 0} + z_-))$ is monotone decreasing with t and $\lim_{t \rightarrow -\infty} \text{YMH}(e^u \cdot (x + z_{\geq 0} + z_-)) = \text{YMH}(x)$, then $\text{YMH}(e^u \cdot (x + z_{\geq 0} + z_-)) \leq \text{YMH}(x)$, and so the above equation implies that

$$|\text{YMH}(y_t) - \text{YMH}(x)| \leq K'_1 e^{K'_2 t}$$

for positive constants $K'_1 = \varepsilon \sqrt{K_1}$ and $K'_2 = \frac{1}{2} K_2$. \square

Lemma 2.25. *Let y_t be a solution to the YMH flow such that $y_t \rightarrow x$ as $t \rightarrow -\infty$. Then for each positive integer k there exists a constant C and a constant $\tau_0 \in \mathbb{R}$ such that*

$$\|y_\tau - x\|_{L_k^2} \leq C \int_{-\infty}^{\tau} \|\text{grad YMH}(y_s)\|_{L^2} ds$$

for all $\tau \leq \tau_0$.

Proof. Recall the interior estimate from [41, Lem. 7.3], [51, Prop. 3.6] which says that for all positive integers k there exists a neighbourhood U of x in the L_k^2 topology and a constant C such that if $y_t \in U$ for all $t \in [0, T]$ then

$$\int_1^T \|\text{grad YMH}(y_t)\|_{L_k^2} dt \leq C \int_0^T \|\text{grad YMH}(y_t)\|_{L^2} dt.$$

The constant C is uniform as long as the Yang-Mills-Higgs functional is uniformly bounded along y_t , the initial condition satisfies a uniform bound on the derivatives of the curvature of the underlying holomorphic bundle with Hermitian metric and the flow line y_t remains in the fixed neighbourhood U of the critical point x (cf. [41, Prop. A]). In particular, the estimates of [51, Lem. 3.14, Cor 3.16] show that this bound on the curvature is satisfied for any initial condition along a given flow line y_t . *A priori* the constant depends on T , however it can be made uniform in T using the following argument. Let C be the constant for $T = 2$. For any $T \geq 2$, let N be an integer greater than T such that $y_t \in U$ for all $t \in [0, N]$. We then have

$$\begin{aligned} \int_1^T \|\text{grad YMH}(y_t)\|_{L_k^2} dt &\leq \sum_{n=1}^{N-1} \int_n^{n+1} \|\text{grad YMH}(y_t)\|_{L_k^2} dt \\ &\leq C \sum_{n=1}^{N-1} \int_{n-1}^{n+1} \|\text{grad YMH}(y_t)\|_{L^2} dt \\ &\leq 2C \int_0^N \|\text{grad YMH}(y_t)\|_{L^2} dt \end{aligned}$$

Since $\lim_{t \rightarrow -\infty} y_t = x$ in the C^∞ topology, then for any $\varepsilon > 0$ there exists τ_0 such that $\tau \leq \tau_0$ implies that $\|y_t - x\|_{L_k^2} < \varepsilon$ for all $t \leq \tau$ and therefore by choosing ε small we can apply the above interior estimate on any interval $[t, \tau]$ for $\tau \leq \tau_0$. Therefore we have the bound

$$\int_t^\tau \|\text{grad YMH}(y_s)\|_{L_k^2} ds \leq 2C \int_{-\infty}^\tau \|\text{grad YMH}(y_s)\|_{L^2} ds$$

For fixed τ the right-hand side of the above inequality is constant, and so

$$\|y_\tau - x\|_{L_k^2} \leq \int_{-\infty}^\tau \|\text{grad YMH}(y_s)\|_{L_k^2} ds \leq 2C \int_{-\infty}^\tau \|\text{grad YMH}(y_s)\|_{L^2} ds$$

□

Proof of Proposition 2.21. After possibly shrinking the neighbourhood U from the previous lemma, we can use the Lojasiewicz inequality (cf. [51, Prop. 3.5]) which implies that for any $y \in U$ there exist constants $C > 0$ and $\theta \in (0, \frac{1}{2}]$ such that

$$(2.19) \quad \|\text{grad YMH}(y)\| \geq C |\text{YMH}(x) - \text{YMH}(y)|^{1-\theta}$$

Along the gradient flow we have the inequality

$$\begin{aligned} \frac{\partial}{\partial t} (\text{YMH}(x) - \text{YMH}(y_t))^\theta &= -\theta (\text{YMH}(x) - \text{YMH}(y_t))^{\theta-1} \frac{\partial}{\partial t} \text{YMH}(y_t) \\ &= \theta (\text{YMH}(x) - \text{YMH}(y_t))^{\theta-1} \|\text{grad YMH}(y_t)\|^2 \\ &\geq C\theta \|\text{grad YMH}(y_t)\| \end{aligned}$$

where the last step follows from the Lojasiewicz inequality (2.19). Since $y_t \rightarrow x$ as $t \rightarrow -\infty$, then integrating the above inequality from $-\infty$ to τ gives us

$$(\text{YMH}(x) - \text{YMH}(y_\tau))^\theta \geq C\theta \int_{-\infty}^{\tau} \|\text{grad YMH}(y_t)\| dt,$$

which is equivalent to

$$\int_{-\infty}^{\tau} \|\text{grad YMH}(y_t)\| dt \leq \frac{1}{C\theta} (\text{YMH}(x) - \text{YMH}(y_\tau))^\theta.$$

Lemma 2.24 shows that

$$(\text{YMH}(x) - \text{YMH}(y_\tau))^\theta \leq (K'_1)^\theta e^{\theta K'_2 t}$$

for all $t \leq 0$. These two estimates together with the result of Lemma 2.25 show that

$$\|y_t - x\|_{L_k^2} \leq C_1 e^{\eta t}$$

for some positive constants C_1, η and all $t \leq 0$. □

3. THE ISOMORPHISM CLASSES IN THE UNSTABLE SET

Given a critical point $x \in \mathcal{B}$, in this section we show that for each $y \in S_x^-$ there exists a smooth gauge transformation $g \in \mathcal{G}^{\mathbb{C}}$ such that $g \cdot y \in W_x^-$ (Proposition 3.14), and conversely for each $y \in W_x^-$ there exists $g \in \mathcal{G}^{\mathbb{C}}$ such that $g \cdot y \in S_x^-$ (Proposition 3.18). As a consequence, the isomorphism classes in the unstable set are in bijective correspondence with the isomorphism classes in the negative slice, and so we have a complete description of these isomorphism classes by Lemma 2.10. This leads to Theorem 3.19 which gives an algebraic criterion for two points to be connected by a flow line.

3.1. Convergence of the scattering construction. The goal of this section is to prove Proposition 3.14, which shows that every point in the negative slice S_x^- is complex gauge equivalent to a point in the unstable set W_x^- .

The construction involves flowing up towards the critical point on the slice using the linearisation of the YMH flow and then flowing down using the YMH flow. A similar idea is used by Hubbard in [19] for analytic flows around a critical point in \mathbb{C}^n , where the flow on the slice is defined by projecting the flow from the ambient space. Hubbard's construction uses the fact that the ambient space is a manifold to (a) define this projection to the negative slice, and (b) define local coordinates in which the nonlinear part of the gradient flow satisfies certain estimates in terms of the eigenvalues for the linearised flow [19, Prop. 4], which is necessary to prove convergence. This idea originated in the study of the existence of scattering states in classical and quantum mechanics. In the context of this paper, one can think of the linearised flow and the YMH flow as two dynamical systems and the goal is to compare their behaviour as $t \rightarrow -\infty$ (see [42, Ch. XI.1] for an overview). As noted in [19], [40] and [42], the eigenvalues of the linearised flow play an important role in comparing the two flows.

The method of this section circumvents the need for a local manifold structure by defining the flow on the slice using the linearised flow and then using the distance-decreasing property of the flow on the space of metrics from [8], [45] (cf. Lemma 2.18) in place of the estimate of [19, Prop. 4] on the nonlinear part of the flow. The entire construction is done in terms of the complex gauge group, and so it is valid

on any subset preserved by $\mathcal{G}^{\mathbb{C}}$, thus avoiding any problems associated with the singularities in the space of Higgs bundles. Moreover, using this method it follows naturally from the Lojasiewicz inequality and the smoothing properties of the heat equation that the backwards YMH flow with initial condition in the unstable set converges in the C^∞ topology.

3.1.1. *A C^0 bound on the metric.* First we derive an *a priori* estimate on the change of metric along the flow. Fix an initial condition $y_0 \in S_x^-$ and let $\beta = \mu(x) = \Lambda(F_A + [\phi, \phi^*]) \in \Omega^0(\text{ad}(E)) \cong \text{Lie}(\mathcal{G})$. In this section we also use the function $\mu_h(y) = \text{Ad}_{g^{-1}}(\mu(g \cdot y))$ from (2.15). The linearised flow with initial condition y_0 has the form $e^{-i\beta t} \cdot y_0$, and the YMH flow (2.3) has the form $g_t \cdot y_0$. Let $f_t = g_t \cdot e^{i\beta t}$ and define $h_t = f_t^* f_t \in \mathcal{G}^{\mathbb{C}}/\mathcal{G}$. This is summarised in the diagram below.




FIGURE 1. Comparison of the gradient flow and the linearised flow.

Lemma 3.1. *For any initial condition $y_0 \in S_x^-$, the induced flow on $\mathcal{G}^{\mathbb{C}}/\mathcal{G}$ satisfies*

$$\frac{dh_t}{dt} = -2ih_t \mu_h(e^{-i\beta t} \cdot y_0) + i\beta h_t + h_t(i\beta)$$

Proof. First compute

$$(3.1) \quad \frac{df_t}{dt} f_t^{-1} = \frac{dg_t}{dt} g_t^{-1} + g_t(i\beta) e^{i\beta t} f_t^{-1} = -i\mu(g_t \cdot y_0) + f_t(i\beta) f_t^{-1}.$$

Then

$$\begin{aligned} \frac{dh_t}{dt} &= \frac{df_t^*}{dt} f_t + f_t^* \frac{df_t}{dt} \\ &= f_t^* \left(\frac{df_t}{dt} f_t^{-1} \right)^* f_t + f_t^* \left(\frac{df_t}{dt} f_t^{-1} \right) f_t \\ &= -f_t^* i\mu(g_t \cdot y_0) f_t + i\beta h_t - f_t^* i\mu(g_t \cdot y_0) f_t + h_t(i\beta) \\ &= -2f_t^* i\mu(g_t \cdot y_0) f_t + i\beta h_t + h_t(i\beta) \\ &= -2ih_t \text{Ad}_{f_t^{-1}}(\mu(g_t \cdot y_0)) + i\beta h_t + h_t(i\beta) \\ &= -2ih_t \mu_h(e^{-i\beta t} \cdot y_0) + i\beta h_t + h_t(i\beta) \end{aligned}$$

where the last step follows from the definition of μ_h in (2.15) and the fact that $e^{-i\beta t} = f_t^{-1} \cdot g_t$. \square

The next estimate gives a bound for $\sup_X \sigma(h_t)$ in terms of $\|y_0 - x\|_{C^0}$.

Lemma 3.2. *For every $\varepsilon > 0$ there exists a constant $C > 0$ such that for any initial condition $y_0 \in S_x^-$ with $\|e^{-i\beta T} \cdot y_0 - x\|_{C^0} < \varepsilon$ we have the estimate $\sup_X \sigma(h_t) \leq C \|e^{-i\beta T} \cdot y_0 - x\|_{C^0}^2$ for all $0 \leq t \leq T$.*

Proof. Taking the trace of the result of Lemma 3.1 gives us

$$\begin{aligned} \frac{d}{dt} \operatorname{Tr} h_t &= \operatorname{Tr} \left(\frac{dh}{dt} \right) = -2i \operatorname{Tr} \left((\mu_h(e^{-i\beta t} \cdot y_0) - \beta) h_t \right) \\ \frac{d}{dt} \operatorname{Tr} h_t^{-1} &= - \operatorname{Tr} \left(h_t^{-1} \frac{dh}{dt} h_t^{-1} \right) = 2i \operatorname{Tr} \left(h_t^{-1} (\mu_h(e^{-i\beta t} \cdot y_0) - \beta) \right) \end{aligned}$$

Therefore

$$\frac{d}{dt} \operatorname{Tr}(h_t) = -2i \operatorname{Tr} \left((\mu_h(e^{-i\beta t} \cdot y_0) - \mu(e^{-i\beta t} \cdot y_0)) h_t \right) - 2i \operatorname{Tr} \left((\mu(e^{-i\beta t} \cdot y_0) - \beta) h_t \right)$$

Lemma 2.20 together with the fact that h_t is positive definite then shows that

$$\begin{aligned} \left(\frac{\partial}{\partial t} + \Delta \right) \operatorname{Tr}(h_t) &\leq -2i \operatorname{Tr} \left((\mu(e^{-i\beta t} \cdot y_0) - \beta) h_t \right) \\ &\leq C_1 \|\mu(e^{-i\beta t} \cdot y_0) - \beta\|_{C^0} \operatorname{Tr}(h_t) \\ &\leq C_1 \|e^{-i\beta t} \cdot y_0 - x\|_{C^0}^2 \operatorname{Tr}(h_t) \quad (\text{by Lemma 2.17}) \end{aligned}$$

A similar calculation shows that

$$\left(\frac{\partial}{\partial t} + \Delta \right) \operatorname{Tr}(h_t^{-1}) \leq C_1 \|e^{-i\beta t} \cdot y_0 - x\|_{C^0}^2 \operatorname{Tr}(h_t^{-1})$$

If we label the eigenvalues of $i\beta$ as $\lambda_1 \leq \dots \leq \lambda_k < 0 \leq \lambda_{k+1} \leq \dots \leq \lambda_n$, then the estimate $\|e^{i\beta s} \cdot (y_0 - x)\|_{C^0}^2 \leq e^{2\lambda_k s} \|y_0 - x\|_{C^0}^2$ from (2.14) gives us

(3.2)

$$\begin{aligned} \left(\frac{\partial}{\partial t} + \Delta \right) \sigma(h_t) &= \left(\frac{\partial}{\partial t} + \Delta \right) (\operatorname{Tr}(h_t) + \operatorname{Tr}(h_t^{-1})) \\ &\leq C_1 \|e^{-i\beta t} \cdot (y_0 - x)\|_{C^0}^2 (\operatorname{Tr}(h_t) + \operatorname{Tr}(h_t^{-1})) \\ &= C_1 \|e^{i\beta(T-t)} \cdot e^{-i\beta T} \cdot (y_0 - x)\|_{C^0}^2 (\operatorname{Tr}(h_t) + \operatorname{Tr}(h_t^{-1})) \\ &\leq C_1 e^{2\lambda_k(T-t)} \|e^{-i\beta T} \cdot (y_0 - x)\|_{C^0}^2 \sigma(h_t) + C_1 e^{2\lambda_k(T-t)} \|e^{-i\beta T} \cdot (y_0 - x)\|_{C^0}^2 \operatorname{rank}(E) \end{aligned}$$

Let $K_1 = C_1 \|e^{-i\beta T} \cdot y_0 - x\|_{C^0}^2$ and $K_2 = C_1 \|e^{-i\beta T} \cdot y_0 - x\|_{C^0}^2 \operatorname{rank}(E)$. Define

$$\nu_t = \sigma(h_t) \exp \left(\frac{K_1}{2\lambda_k} e^{2\lambda_k(T-t)} \right) - \int_0^t K_2 e^{2\lambda_k(T-s)} \exp \left(\frac{K_1}{2\lambda_k} e^{2\lambda_k(T-s)} \right) ds$$

Note that $\nu_0 = 0$ since $h_0 = \operatorname{id}$. A calculation using (3.2) then shows that

$$\left(\frac{\partial}{\partial t} + \Delta \right) \nu_t \leq 0$$

and so $\sup_X \nu_t \leq \sup_X \nu_0 = 0$ by the maximum principle. Therefore

$$\begin{aligned} \sup_X \sigma(h_t) &\leq \exp \left(-\frac{K_1}{2\lambda_k} e^{2\lambda_k(T-t)} \right) \int_0^t K_2 e^{2\lambda_k(T-s)} \exp \left(\frac{K_1}{2\lambda_k} e^{2\lambda_k(T-s)} \right) ds \\ &\leq \exp \left(-\frac{K_1}{2\lambda_k} \right) \int_0^t K_2 e^{2\lambda_k(T-s)} ds \leq C \|e^{-i\beta T} \cdot y_0 - x\|_{C^0}^2 \end{aligned}$$

for some constant C , since $\lambda_k < 0$, $0 \leq s \leq t < T$, K_1 is bounded since $\|e^{-i\beta T} \cdot y_0 - x\|_{C^0} < \varepsilon$ by assumption and K_2 is proportional to $\|e^{-i\beta T} \cdot y_0 - x\|_{C^0}^2$. \square

3.1.2. C^∞ convergence in the space of metrics. Now consider the case of a fixed $y_0 \in S_x^-$ and define $y_t = e^{i\beta t} \cdot y_0$. Given any $y \in S_x^-$, define $g_s(y) \in \mathcal{G}^C$ to be the unique solution of (2.3) such that $g_s(y) \cdot y$ is the solution to the YMH flow at time s with initial condition y , define $f_s(y) = g_s(e^{i\beta s} \cdot y) \cdot e^{i\beta s} \in \mathcal{G}^C$, and define $h_s(y) = f_s(y)^* f_s(y)$ to be the associated change of metric. For any nonnegative s, t , we can use the result of the previous lemma with y_0 and T replaced by $e^{i\beta s} \cdot y_t$ and s respectively to obtain

$$(3.3) \quad \sup_X \sigma(h_s(y_t)) \leq C \|y_t - x\|_{C^0}^2 = C \|e^{i\beta t} \cdot (y_0 - x)\|_{C^0}^2 \leq C e^{2\lambda_k t} \|y_0 - x\|_{C^0}^2.$$

As a special case, given $t_1 > t_2 \geq 0$, setting $t = t_2$ and $s = t_1 - t_2$ in the above estimate gives us

$$(3.4) \quad \sup_X \sigma(h_{t_1-t_2}(y_{t_2})) \leq C \|y_{t_2} - x\|_{C^0}^2.$$

This is summarised in the diagram below.




FIGURE 2. Comparison of $f_{t_1}(y_0) \cdot y_0$ and $f_{t_2}(y_0) \cdot y_0$.

In the above notation, $h_t(y_0) = f_t(y_0)^* f_t(y_0)$ is the change of metric associated to the gauge transformation $f_t(y_0)$ in the above diagram.

Proposition 3.3. $h_t(y_0) \xrightarrow{C^0} h_\infty(y_0) \in \mathcal{G}^C/\mathcal{G}$ as $t \rightarrow \infty$. The limit depends continuously on the initial condition y_0 . The rate of convergence is given by

$$(3.5) \quad \sup_X \sigma(h_t(y_0)(h_\infty(y_0))^{-1}) \leq C_2 e^{2\lambda_k t} \|y_0 - x\|_{C^0}^2$$

where $C_2 > 0$ is a constant depending only on the orbit $\mathcal{G} \cdot x$.

Proof. Let $t_1 > t_2 \geq T$. The estimate (3.4) shows that

$$\sup_X \sigma(h_{t_1-t_2}(y_{t_2})) \leq C \|y_{t_2} - x\|_{C^0}^2 \leq C e^{2\lambda_k t_2} \|y_0 - x\|_{C^0}^2 \leq C e^{2\lambda_k T} \|y_0 - x\|_{C^0}^2.$$

Recall from (2.17) that

$$\sigma(h_{t_1}(y_0)h_{t_2}(y_0)^{-1}) = \sigma((f_{t_1}(y_0)f_{t_2}(y_0)^{-1})^* f_{t_1}(y_0)f_{t_2}(y_0)^{-1}).$$

The distance-decreasing formula of Lemma 2.18 shows that

$$\sup_X \sigma((f_{t_1}(y_0)f_{t_2}(y_0)^{-1})^* f_{t_1}(y_0)f_{t_2}(y_0)^{-1}) \leq \sup_X \sigma(h_{t_1-t_2}(y_{t_2})).$$

Therefore the distance (measured by σ) between the two metrics $h_{t_1}(y_0)$ and $h_{t_2}(y_0)$ satisfies the following bound

$$\begin{aligned} \sup_X \sigma(h_{t_1}(y_0)h_{t_2}(y_0)^{-1}) &= \sup_X \sigma((f_{t_1}(y_0)f_{t_2}(y_0)^{-1})^* f_{t_1}(y_0)f_{t_2}(y_0)^{-1}) \\ &\leq \sup_X \sigma(h_{t_1-t_2}(y_{t_2})) \leq C_2 e^{2\lambda_k T} \|y_0 - x\|_{C^0}^2 \end{aligned}$$

and so $h_t(y_0)$ is a Cauchy sequence in C^0 with a unique limit $h_\infty \in \mathcal{G}^C/\mathcal{G}$. The above equation shows that the rate of convergence is given by (3.5).

Since the finite-time Yang-Mills-Higgs flow and linearised flow both depend continuously on the initial condition, then $h_t(y_0)$ depends continuously on y_0 for each $t > 0$. Continuous dependence of the limit then follows from the estimate (3.5). \square

Now we can improve on the previous estimates to show that $h_t(y_0)$ converges in the smooth topology along a subsequence, and therefore the limit h_∞ is C^∞ . Define $z_t = f_t(y_0) \cdot y_0$, where $y_0 \in S_x^-$ and $f_t(y_0) \in \mathcal{G}^C$ are as defined in the previous proposition. Given a Higgs bundle $z_t = (\bar{\partial}_A, \phi)$, let ∇_A denote the covariant derivative with respect to the metric connection associated to $\bar{\partial}_A$.

Lemma 3.4. *For each initial condition $y_0 \in S_x^-$, there is a uniform bound on $\sup_X |\nabla_A^\ell \mu(z_t)|$ and $\sup_X |\nabla_A^\ell \phi|$ for each $\ell \geq 0$.*


Proof. Since $\{e^{i\beta t} \cdot y_0 : t \in [0, \infty]\}$ is a compact curve in the space of C^∞ Higgs bundles connecting two C^∞ Higgs bundles y_0 and x , then $\sup_X |\mu(e^{i\beta t} \cdot y_0)|$ and $\sup_X |\nabla_A \phi|$ are both uniformly bounded along the sequence $e^{i\beta t} \cdot y_0$. By construction, z_t is the time t YMH flow with initial condition $e^{i\beta t} \cdot y_0$. Along the YMH flow, for each ℓ the quantities $\sup_X |\nabla_A^\ell \mu|$ and $\sup_X |\nabla_A^\ell \phi|$ are both uniformly bounded by a constant depending on the value of $\sup_X |\mu|$ and $\sup_X |\nabla_A \phi|$ at the initial condition (cf. [51, Sec. 3.2]). Since these quantities are uniformly bounded for the initial conditions, then the result follows. \square

Corollary 3.5. *There is a subsequence t_n such that $h_{t_n} \rightarrow h_\infty$ in the C^∞ topology. Therefore h_∞ is C^∞ .*

Proof. Since z_t is contained in the complex gauge orbit of y_0 for all t , then [51, Lem. 3.14] shows that the uniform bound on $|\nabla_A^\ell \mu(z_t)|$ from the previous lemma implies a uniform bound on $|\nabla_A^\ell F_A|$ for all ℓ . Therefore, since Proposition 3.3 shows that h_t converges in C^0 , then the estimates of [8, Lem. 19 & 20] show that h_t is bounded in C^ℓ for all ℓ , and so there is a subsequence h_{t_n} converging in the C^∞ topology. \square

3.1.3. C^∞ convergence in the space of Higgs bundles. In this section we show that the scattering construction converges in the C^∞ topology on the space of Higgs bundles. As a consequence of the methods, we obtain an estimate that shows the solution to the reverse heat flow constructed in Section 3.1.4 converges to the critical point x in the smooth topology.

This section uses a slightly modified version of the flow from the previous section, defined as follows. Given $y_0 \in S_x^-$ and $t > 0$, let $x_s = g_s \cdot e^{i\beta t} \cdot y_0$ be the time s solution to the YMH flow (2.3) with initial condition $e^{i\beta t} \cdot y_0$, let $s(t)$ be the unique point in time such that $\text{YMH}(x_{s(t)}) = \text{YMH}(y_0)$ and define $t' = \min\{t, s(t)\}$. At a critical point, the curvature $i * (F_A + [\phi, \phi^*])$ takes the form of a diagonal matrix whose entries are rational numbers with denominator bounded above by $\text{rank}(E)$ (cf. (2.6)), and therefore the critical values of YMH are discrete, which implies that t' is well-defined for small values of $\text{YMH}(x) - \text{YMH}(y_0)$. The figure below shows the two cases $t' = s(t) \leq t$ and $t' = t < s(t)$.



Now define $z_t = g_{t'} \cdot e^{i\beta t} \cdot y_0$ and $w_t = e^{i\beta(t-t')} \cdot y_0$. Note that $z_t = g_{t'} \cdot e^{i\beta t'} \cdot w_t$ and so the results of the previous section show that the C^0 norm of the change of metric connecting w_t and z_t is bounded. Therefore Corollary 2.12 shows that w_t and z_t are both uniformly bounded away from x .

Lemma 3.6. *There exists $T > 0$ such that $t - t' \leq T$ for all t .*

Proof. If $s(t) \geq t$ then $t' = t$ and the desired inequality holds. Therefore the only non-trivial case is $s(t) < t$. Since $\text{YMH}(z_t) = \text{YMH}(y_0)$ and YMH is continuous in the L_1^2 norm on \mathcal{B} , then there exists a neighbourhood V of x such that $z_t \notin V$ for all t . We also have $z_t = f_{t'} \cdot w_t$ with $f_{t'} = g_{t'} e^{i\beta t'}$ such that $h_t = f_{t'}^* f_{t'}$ satisfies $\sup_X \sigma(h_t) \leq C \|w_t - x\|_{C^0}^2 \leq C \|y_0 - x\|_{C^0}^2$ by Lemma 3.2, and so Corollary 2.12 shows that there exists a neighbourhood U of x in the L_1^2 topology on \mathcal{B} such that $w_t \notin U$. Therefore there exists $\eta > 0$ such that $\|w_t - x\|_{L_1^2} \geq \eta$ and $\|z_t - x\|_{L_1^2} \geq \eta$.

Since $w_t = e^{i\beta(t-t')} \cdot y_0$ and $e^{i\beta s} \cdot y_0$ converges to x as $s \rightarrow \infty$, then there exists T such that $t - t' \leq T$ for all t , since otherwise $\|w_t - x\|_{L_1^2} < \eta$ for some t which contradicts the inequality from the previous paragraph. \square

Remark 3.7. Since $t - T \leq t' \leq s(t)$, then $\text{YMH}(g_{t-T} \cdot e^{i\beta t} \cdot y_0) \geq \text{YMH}(y_0)$ for all t . Equivalently, $\text{YMH}(g_{t-T} \cdot e^{i\beta(t-T)} \cdot (e^{i\beta T} \cdot y_0)) \geq \text{YMH}(y_0)$. Therefore, by using the fixed initial condition $e^{i\beta T} \cdot y_0$ we can guarantee that the YMH functional remains greater than $\text{YMH}(y_0)$. This is used in Proposition 3.9 (see also Remark 3.12).

Next we use the Lojasiewicz inequality to derive a uniform bound on $\|z_t - x\|_{L_1^2}$. The original Lojasiewicz inequality for a real analytic manifold M and a critical point $c \in M$ of an analytic function $f : M \rightarrow \mathbb{R}$ gives a neighbourhood of c on which $\text{grad } f(z)$ is bounded below by $|f(z) - f(c)|^r$ for some $r > 0$ (cf. [33]). Simon [44] extended this to an infinite-dimensional setting and used this to prove convergence of certain nonlinear evolution equations. This was further extended by Rade [41] who proved that the Yang-Mills flow converges when the base manifold has dimension 2 or 3. In the next lemma we use the inequality in a different way to show that the nonlinear flow z_t remains in a given neighbourhood of the critical point x if the linearised flow remains in a small enough neighbourhood of x .

Lemma 3.8. *Given $\varepsilon > 0$ there exists $\delta > 0$ such that for each $y_0 \in S_x^-$ with $\|y_0 - x\|_{L_1^2} < \delta$ there exists a neighbourhood U of x in the L_1^2 topology such that $\|z_t - x\|_{L_1^2} < \varepsilon$ for all t such that $e^{i\beta t} \cdot y_0 \in U$.*

Proof. Recall from [51, Prop. 3.5] that there exists $\varepsilon_1 > 0$ and constants $C > 0$ and $\theta \in (0, \frac{1}{2})$ such that the Lojasiewicz inequality

$$(3.6) \quad \|\text{grad YM}(z)\|_{L^2} \geq C |\text{YM}(x) - \text{YM}(z)|^{1-\theta}$$

holds for all z such that $\|z - x\|_{L_1^2} < \varepsilon_1$. Recall the interior estimate [51, Prop. 3.6] which says that for any positive integer k there exists $\varepsilon_2 > 0$ and a constant C'_k such that for any solution $x_s = g_s \cdot e^{i\beta t} \cdot y_0$ to the YM flow with initial condition $e^{i\beta t} \cdot y_0$ which satisfies $\|x_s - x\|_{L_k^2} < \varepsilon_2$ for all $0 \leq s \leq S$, then we have

$$(3.7) \quad \int_1^S \|\text{grad YM}(x_s)\|_{L_k^2} dt \leq C'_k \int_0^S \|\text{grad YM}(x_s)\|_{L^2} dt.$$

where the constant C'_k is uniform over all initial conditions in a given $\mathcal{G}^{\mathbb{C}}$ orbit and for all S such that $\|x_s - x\|_{L_k^2} < \varepsilon_2$ for all $s \in [0, S]$ (cf. Lemma 2.25). Define $\varepsilon' = \min\{\varepsilon, \varepsilon_1, \varepsilon_2\}$. A calculation using (3.6) (cf. [44]) shows that any flow line x_s which satisfies $\|x_s - x\|_{L_1^2} < \varepsilon'$ for all $s \in [0, t']$ also satisfies the gradient estimate

$$C\theta \|\text{grad YM}(x_s)\|_{L^2} \leq \frac{\partial}{\partial s} |\text{YM}(x) - \text{YM}(x_s)|^\theta$$

and so if $\|x_s - x\|_{L_1^2} < \varepsilon'$ for all $s < t'$ then

$$(3.8) \quad \int_0^{t'} \|\text{grad YM}(x_s)\|_{L^2} ds \leq \frac{1}{C\theta} \left(|\text{YM}(x) - \text{YM}(x_{t'})|^\theta - |\text{YM}(x) - \text{YM}(x_0)|^\theta \right) \\ \leq \frac{1}{C\theta} |\text{YM}(x) - \text{YM}(x_{t'})|^\theta$$

Let $k = 1$ in (3.7) and choose $\delta > 0$ so that $\|y_0 - x\|_{L_1^2} < \delta$ implies that $\frac{1}{C\theta} |\text{YM}(x) - \text{YM}(y_0)|^\theta \leq \frac{\varepsilon'}{3C'_1}$, where C and θ are the constants from the Lojasiewicz inequality (3.6) and C'_1 is the constant from (3.7) for $k = 1$. Therefore, since $\text{YM}(y_0) = \text{YM}(x_{t'}) < \text{YM}(x_\tau) \leq \text{YM}(x)$ for all $\tau < t'$, then

$$(3.9) \quad \frac{1}{C\theta} |\text{YM}(x) - \text{YM}(x_\tau)|^\theta \leq \frac{\varepsilon'}{3C'_1} \quad \text{for all } \tau < t'.$$

Since the finite-time YM flow depends continuously on the initial condition in the L_1^2 norm by [51, Prop. 3.4], then there exists a neighbourhood U of x such that $x_0 \in U$ implies that $\|x_1 - x\|_{L_1^2} < \frac{1}{3}\varepsilon'$. Choose t

large so that $e^{i\beta t} \cdot y_0 = e^{i\beta t'} \cdot w_t \in U$ and let $x_s = g_s \cdot e^{i\beta t} \cdot y_0$ be the solution to the YMH flow at time s with initial condition $x_0 = e^{i\beta t} \cdot y_0$. Note that $x_{t'} = z_t$. Define

$$\tau = \sup\{s \mid \|x_r - x\|_{L_1^2} < \varepsilon' \text{ for all } r \leq s\}$$

and note that $\tau > 0$. By definition of τ , the Lojasiewicz inequality (3.6) and the interior estimate (3.8) are valid for the flow line x_s on the interval $[0, \tau]$. If $\tau < t'$, then (3.8) and (3.9) imply that

$$\begin{aligned} \|x_\tau - x\|_{L_1^2} &\leq \|x_1 - x\|_{L_1^2} + \|x_\tau - x_1\|_{L_1^2} \\ &< \frac{1}{3}\varepsilon' + \int_1^\tau \|\text{grad YMH}(x_s)\|_{L_1^2} ds \\ &\leq \frac{1}{3}\varepsilon' + C_1' \int_0^\tau \|\text{grad YMH}(x_s)\|_{L_2} ds \\ &\leq \frac{1}{3}\varepsilon' + \frac{C_1'}{C\theta} |\text{YMH}(x) - \text{YMH}(x_\tau)|^\theta \leq \frac{1}{3}\varepsilon' + \frac{1}{3}\varepsilon' \end{aligned}$$

contradicting the definition of τ as the supremum. Therefore $t' \leq \tau$ and the same argument as above shows that $\|x_{t'} - x_0\|_{L_1^2} < \frac{2}{3}\varepsilon'$, so we conclude that $z_t = x_{t'}$ satisfies $\|z_t - x\|_{L_1^2} < \frac{2}{3}\varepsilon' < \varepsilon$ for all t such that $e^{i\beta t} \cdot y_0 \in U$. \square

Now that we have a uniform L_1^2 bound on $z_t - x$, then we can apply the same idea using the interior estimate (3.7) as well as continuous dependence on the initial condition in the L_k^2 norm from [51, Prop. 3.4] to prove the following uniform L_k^2 bound on $z_t - x$.

Lemma 3.9. *Given $\varepsilon > 0$ and a positive integer k there exists $\delta > 0$ such that if $\|y_0 - x\|_{L_1^2} < \delta$ then there exists a neighbourhood U of x in the L_k^2 topology such that $\|z_t - x\|_{L_k^2} < \varepsilon$ for all t such that $e^{i\beta t} \cdot y_0 \in U$.*

Now we can prove that there is a limit z_∞ in the space of C^∞ Higgs bundles. In Section 3.1.4 we will show that $z_\infty \in W_x^-$.

Proposition 3.10. *For each $y_0 \in S_x^-$, let z_t be the sequence defined above. Then there exists $z_\infty \in \mathcal{B}$ such that for each positive integer k there exists a subsequence of z_t converging to z_∞ strongly in L_k^2 .*

Proof. The previous estimate with $k = 2$ shows that $\|z_t - x\|_{L_2^2}$ is bounded. Compactness of the embedding $L_{k+1}^2 \hookrightarrow L_k^2$ shows that there is a subsequence $\{z_{t_n}\}$ converging strongly to a limit z_∞ in L_1^2 .

For any $k > 1$, the same argument applied to the subsequence $\{z_{t_n}\}$ from the previous paragraph shows that there exists a further subsequence, which we denote by $\{z_{t_{n_j}}\}$, which converges strongly in L_k^2 . Since $z_{t_{n_j}} \xrightarrow{L_1^2} z_\infty$ then the limit in L_k^2 of $z_{t_{n_j}}$ must be z_∞ also. Therefore z_∞ is a C^∞ Higgs pair. \square

Finally, we can prove that z_∞ is gauge-equivalent to y_0 . Recall the constant T from Lemma 3.6 and let $\varphi(z_t, s)$ denote the time s downwards YMH flow (2.4) with initial condition z_t . The gauge transformation $f_t(y_0) \in \mathcal{G}^C$ from Proposition 3.3 satisfies $f_t(y_0) \cdot y_0 = \phi(z_t, t - t')$.

For any k , let z_{t_n} be a subsequence converging strongly to z_∞ in L_k^2 . Such a subsequence exists by Proposition 3.10. Since $0 \leq t_n - t'_n \leq T$ for all n then there exists $s \in [0, T]$ and a subsequence $\{t_{n_\ell}\}$ such that $t_{n_\ell} - t'_{n_\ell} \rightarrow s$. Since the finite-time YMH flow depends continuously on the initial condition in L_k^2 , then $f_{t_{n_\ell}}(y_0) \cdot y_0 = \varphi(z_{t_{n_\ell}}, t_{n_\ell} - t'_{n_\ell})$ converges to $z_\infty^0 := \varphi(z_\infty, s)$ strongly in L_k^2 . After taking a

further subsequence if necessary, the method of Section 3.1.2 shows that the change of metric associated to $f_{t_{n_\ell}}(y_0)$ converges strongly in L_{k+1}^2 . Therefore, since the action of the Sobolev completion $\mathcal{G}_{L_{k+1}^2}^{\mathbb{C}}$ on $\mathcal{B}_{L_k^2}$ is continuous, then $\varphi(z_\infty, s)$ (and hence z_∞) is related to y_0 by a gauge transformation in $\mathcal{G}_{L_{k+1}^2}^{\mathbb{C}}$. Since y_0 and z_∞ are both smooth Higgs pairs then an elliptic regularity argument shows that this gauge transformation is smooth. Therefore we have proved the following result.

Proposition 3.11. *Given any $y_0 \in S_x^-$, let z_∞ be the limit from Proposition 3.10. Then there exists a smooth gauge transformation $g \in \mathcal{G}^{\mathbb{C}}$ such that $z_\infty = g \cdot y_0$.*

Since $z_\infty^0 = \varphi(z_\infty, s)$ is related to z_∞ by the finite-time flow and s is bounded, then we have the following estimate for $\|z_\infty^0 - x\|_{L_k^2}$. Note that this requires a bound on $\|y_0 - x\|_{L_1^2}$ for the estimates of this section to work, and a bound on $\|y_0 - x\|_{C^0}$ for the estimates of Lemma 3.2 to work.

Corollary 3.12. *For all $\varepsilon > 0$ there exists $\delta > 0$ such that $\|y_0 - x\|_{L_1^2} + \|y_0 - x\|_{C^0} < \delta$ implies $\|z_\infty^0 - x\|_{L_k^2} < \varepsilon$.*

Remark 3.13. The previous proof uses the fact that the *finite-time* flow depends continuously on the initial condition. The limit of the downwards YMH flow as $t \rightarrow \infty$ depends continuously on initial conditions within the same Morse stratum (cf. [51, Thm. 3.1]). It is essential that the constant T from Lemma 3.6 is finite (which follows from Corollary 2.12) in order to guarantee that z_∞ and $\varphi(z_\infty, s)$ are gauge equivalent. Without a bound on T , it is possible that z_∞ may be in a different Morse stratum to $\lim_{t \rightarrow \infty} \varphi(z_t, t - t')$.

3.1.4. *Constructing a convergent solution to the backwards YMH flow.* In this section we show that the limit z_∞ is in the unstable set W_x^- .

Proposition 3.14. *For each $y_0 \in S_x^-$ there exists $g \in \mathcal{G}^{\mathbb{C}}$ such that $g \cdot y_0 \in W_x^-$.*

Proof. In what follows, fix any positive integer k . Given $y_0 \in S_x^-$, let $z_t^0 = f_t(y_0) \cdot y_0$, where f_t is the complex gauge transformation from Proposition 3.3. Again using $\varphi(z, s)$ to denote the time s downwards YMH flow with initial condition z , Proposition 3.11 shows that there exists $z_\infty^0 := \varphi(z_\infty, s)$ and a subsequence $\{z_{t_n}^0\}$ such that $z_{t_n}^0 \rightarrow z_\infty^0$ strongly in L_k^2 .

Given y_0 , define z_∞^0 as above, and now consider any $s > 0$. Let $y_s = e^{i\beta s} \cdot y_0$ and define $z_t^{-s} = f_t(y_s) \cdot y_s$. By definition, $z_t^0 = \varphi(z_{t-s}^{-s}, s)$, i.e. z_t^0 is the downwards YMH flow for time s with initial condition z_{t-s}^{-s} . Applying Proposition 3.10 to the subsequence $\{z_{t_n-s}^{-s}\}$ shows that there is a subsequence $\{z_{t_{n_j}-s}^{-s}\}$ converging in L_k^2 to some z_∞^{-s} . Since the YMH flow for finite time s depends continuously on the initial condition (cf. [51, Prop. 3.4]) then $z_{t_{n_j}}^0 = \varphi(z_{t_{n_j}-s}^{-s}, s)$ converges to $\varphi(z_\infty^{-s}, s)$. Since we already have $z_{t_n}^0 \rightarrow z_\infty^0$, then $z_\infty^0 = \varphi(z_\infty^{-s}, s)$.


Therefore, for any $s > 0$ we have constructed a solution to the YMH flow on $[-s, 0]$ connecting z_∞^0 and z_∞^{-s} . Proposition A.1 shows that this solution must be unique for each s , and so we can extend this to a solution on the time interval $(-\infty, 0]$.

Moreover, we also have the uniform bound from Corollary 3.12 which shows that for all $\varepsilon > 0$ there exists $\delta > 0$ such that $\|z_\infty^{-s} - x\|_{L_k^2} \leq \varepsilon$ for all y_0 such that $\|y_0 - x\|_{L_1^2} < \delta$. Therefore as $s \rightarrow \infty$, the

sequence z_∞^{-s} converges strongly to x in the L_k^2 norm for any k , and so $z_\infty^0 = g \cdot y_0 \in W_x^-$. Proposition 2.21 then shows that the convergence is exponential in each Sobolev norm. \square

3.2. Convergence of the inverse process. In this section we consider the inverse procedure to that of the previous section and prove that each point in the unstable set W_x^- is gauge equivalent to a point in the negative slice S_x^- . The idea is similar to that of the previous section, except that here we use the modified flow from Section 2.3 instead of the Yang-Mills-Higgs flow (2.4).

3.2.1. A C^0 bound in the space of metrics. Given $y_0 \in W_x^-$, let $y_t = g_t \cdot y_0$ be the solution to the modified flow (2.11) with initial condition y_0 . Define $f_t = g_t \cdot e^{i\beta t}$ and let $h_t = f_t^* f_t$. This is summarised in the diagram below.



Using a similar calculation as the previous section, we have the same expression for the change of metric as in Lemma 3.1.

Lemma 3.15. *For any initial condition $y_0 \in W_x^-$, the induced flow on $\mathcal{G}^{\mathbb{C}}/\mathcal{G}$ satisfies*

$$(3.10) \quad \frac{dh_t}{dt} = -2ih_t\mu_h(e^{-i\beta t} \cdot y_0) + i\beta h_t + ih_t\beta.$$

Proof. A similar calculation as in the proof of Lemma 3.1 (this time using the modified flow (2.11)) shows that

$$\frac{df_t}{dt} f_t^{-1} = -i\mu(g_t \cdot y_0) + \gamma(g_t \cdot y_0) + f_t(i\beta)f_t^{-1}.$$

Then

$$\begin{aligned} \frac{dh_t}{dt} &= f_t^* \left(\frac{df}{dt} f_t^{-1} \right)^* f_t + f_t^* \left(\frac{df}{dt} f_t^{-1} \right) f_t \\ &= f_t^* (-i\mu(g_t \cdot y_0) - \gamma(g_t \cdot y_0) + (f_t^*)^{-1}(i\beta)f_t^* - i\mu(g_t \cdot y_0) + \gamma(g_t \cdot y_0) + f_t(i\beta)f_t^{-1}) f_t \\ &= -2ih_t f_t^{-1} \mu(g_t \cdot y_0) f_t + i\beta h_t + ih_t \beta \\ &= -2ih_t \mu_h(e^{-i\beta t} \cdot y_0) + i\beta h_t + ih_t \beta. \end{aligned} \quad \square$$

Lemma 3.16. *For every $\varepsilon > 0$ there exists a constant $C > 0$ such that for any initial condition $y_0 \in W_x^-$ with $\|e^{-i\beta T} \cdot y_0 - x\|_{C^1} + \|g_T \cdot y_0 - x\|_{C^1} < \varepsilon$ we have the estimate*

$$\sup_X \sigma(h_t) \leq C \left(\|e^{-i\beta T} \cdot y_0 - x\|_{C^1} + \|g_T \cdot y_0 - x\|_{C^1} \right)$$

for all $0 \leq t \leq T$.

Proof. In contrast to the proof of Lemma 3.2, $e^{-i\beta t} \cdot y_0$ is not in the slice S_x and so it satisfies the inequality $\|\mu(e^{-i\beta t} \cdot y_0) - \beta\|_{C^0} \leq C' \|e^{-i\beta t} \cdot y_0 - x\|_{C^1}$ instead of the quadratic bound of Lemma 2.17. Using this inequality, the same idea as in the proof of Lemma 3.2 leads to the bound

$$(3.11) \quad \begin{aligned} \left(\frac{\partial}{\partial t} + \Delta\right) \sigma(h_t) &\leq C_1 \|e^{-i\beta t} \cdot (y_0 - x)\|_{C^1} (\text{Tr}(h_t) + \text{Tr}(h_t^{-1})) \\ &= C_1 \|e^{-i\beta t} \cdot (y_0 - x)\|_{C^1} \sigma(h_t) + 2C_1 \|e^{-i\beta t} \cdot (y_0 - x)\|_{C^1} \text{rank}(E) \end{aligned}$$

In general, if the heat operator is bounded for all $t \geq 0$

$$\left(\frac{\partial}{\partial t} + \Delta\right) f(p, t) \leq C(t) f(p, t) + D(t), \quad p \in X, t \in [0, \infty)$$

for some nonnegative functions $C(t)$ and $D(t)$ independent of $p \in X$, then $f(p, t)$ satisfies the bound

$$(3.12) \quad f(p, t) \leq \exp\left(\int_0^t C(s) ds\right) \left(\int_0^t D(s) ds + f(p, 0)\right)$$

Therefore, since $h_0 = \text{id}$ and so $\sigma(h_0) = 0$, then (3.11) implies that the problem reduces to finding a bound for $\int_0^t \|e^{-i\beta s} \cdot (y_0 - x)\|_{C^1} ds$. Proposition 2.21 shows that the backwards flow with initial condition in W_x^- converges exponentially to x in every Sobolev norm. Therefore there exists a neighbourhood U of x such that if $g_T \cdot y_0 \in U$ then there exist positive constants C_1 and η such that the following estimate holds

$$\|y_0 - x\|_{C^1} \leq C_1 e^{-\eta T} \|g_T \cdot y_0 - x\|_{C^1}.$$

Recall the eigenbundles $\text{End}(E)_-$, $\text{End}(E)_0$ and $\text{End}(E)_+$ from Section 2. The above estimate shows that each component of $y_0 - x$ in $\text{End}(E)_-$, $\text{End}(E)_0$ and $\text{End}(E)_+$ is bounded by $C_1 e^{-\eta T} \|g_T \cdot y_0 - x\|_{C^1}$. Since the component of $e^{-i\beta t} \cdot (y_0 - x)$ in $\text{End}(E)_+$ is exponentially decreasing with t then

$$\int_0^T \|(e^{-i\beta t} \cdot y_0 - x)_{\text{End}(E)_+}\|_{C^1} dt \leq C'_1 \|y_0 - x\|_{C^1} \leq C_1 e^{-\eta T} \|g_T \cdot y_0 - x\|_{C^1}.$$

The component of $e^{-i\beta t} \cdot (y_0 - x)$ in $\text{End}(E)_0$ is constant with respect to t , and so

$$\int_0^T \|(e^{-i\beta t} \cdot y_0 - x)_{\text{End}(E)_0}\|_{C^1} dt \leq C'_2 T \|y_0 - x\|_{C^1} \leq C_2 T e^{-\eta T} \|g_T \cdot y_0 - x\|_{C^1}.$$

Finally, the component of $e^{-i\beta t} \cdot (y_0 - x)$ in $\text{End}(E)_-$ is exponentially increasing. In general, for any $\lambda > 0$ we have the estimate

$$\int_0^T e^{\lambda t} dt = \frac{1}{\lambda} (e^{\lambda T} - 1) \leq \frac{1}{\lambda} e^{\lambda T}$$

and so we have the bound

$$\int_0^T \|(e^{-i\beta t} \cdot y_0 - x)_{\text{End}(E)_-}\|_{C^1} dt \leq C_3 \|e^{-i\beta T} \cdot (y_0 - x)\|_{C^1}.$$

Combining the estimates for the three components shows that the integral

$$I(t) = \int_0^t \|e^{-i\beta s} \cdot (y_0 - x)\|_{C^1} ds$$


is bounded by

$$I(t) \leq C_1 e^{-\eta T} \|g_T \cdot y_0 - x\|_{C^1} + C_2 T e^{-\eta T} \|g_T \cdot y_0 - x\|_{C^1} + C_3 \|e^{-i\beta T} \cdot y_0 - x\|_{C^1}$$

The inequality (3.12) together with the assumption $\|g_T \cdot y_0 - x\|_{C^1} + \|e^{-i\beta T} \cdot y_0 - x\|_{C^1} < \varepsilon$ shows that there exists a constant C such that

$$\sup_X \sigma(h_t) \leq C \left(\|e^{-i\beta T} \cdot y_0 - x\|_{C^1} + \|g_T \cdot y_0 - x\|_{C^1} \right) \quad \square$$

3.2.2. Convergence in the space of Higgs bundles. In this section we use a method analogous to that of Section 3.1.3 to show that the sequence converges in the space of Higgs bundles and that the limit is gauge equivalent to y_0 . Given $y_0 \in W_x^-$ and $t \in (-\infty, 0]$, define $s < 0$ by $\|e^{i\beta s} \cdot g_t(y_0) \cdot y_0 - x\|_{L_k^2} = \|y_0 - x\|_{L_k^2}$. Note that this is well-defined for small values of $\|y_0 - x\|_{L_k^2}$ since Lemma 2.16 shows that $g_t(y_0) \cdot y_0 \rightarrow x$ in the C^∞ topology as $t \rightarrow -\infty$ and for $s < 0$ the action of $e^{i\beta s}$ exponentially increases the C^0 norm of the component of $g_t(y_0) \cdot y_0$ in $\text{End}(E)_-$. Now define $t' := \max\{t, s\} < 0$, let $f_t(y_0) = e^{i\beta t'} \cdot g_{t-t'}(y_0) \cdot y_0$ and $z_t := e^{i\beta t'} \cdot g_t(y_0) \cdot y_0 = f_t(y_0) \cdot g_{t-t'}(y_0) \cdot y_0$. Let $h_t = f_t^* f_t$ be the associated change of metric.



Lemma 3.16 then shows that $\sup_X \sigma(h_t) \leq C (\|z_t - x\|_{C^1} + \|g_{t-t'}(y_0) \cdot y_0 - x\|_{C^1})$. Since either $\|z_t - x\|_{L_k^2} = \|y_0 - x\|_{L_k^2}$ (when $t < t'$) or $g_{t-t'}(y_0) \cdot y_0 = y_0$ (when $t' = t$), then Corollary 2.12 shows that $g_{t-t'}(y_0) \cdot y_0$ and z_t are both bounded away from x in the L_k^2 norm. As a consequence, $|t - t'|$ is uniformly bounded in the same way as Lemma 3.6. Therefore

$$(3.13) \quad \|e^{i\beta t} \cdot g_t(y_0) \cdot y_0 - x\|_{L_k^2} = \|e^{i\beta(t-t')} \cdot z_t - x\|_{L_k^2} \leq C' \|z_t - x\|_{L_k^2} = C' \|y_0 - x\|_{L_k^2}$$

for some constant C' , which implies that there is a subsequence of $e^{i\beta t} \cdot g_t(y_0) \cdot y_0$ converging strongly to a limit z_∞^0 in L_{k-1}^2 . Since this is true for all k , then z_∞^0 is a C^∞ Higgs pair.

A special case of (3.13) is


$$(3.14) \quad \|e^{i\beta t} \cdot g_t(y_0) \cdot y_0 - x\|_{C^1} \leq C \|e^{i\beta t} \cdot g_t(y_0) \cdot y_0 - x\|_{L_k^2} \leq C' \|y_0 - x\|_{L_k^2}$$

for any k such that $L_k^2 \hookrightarrow C^1$ is an embedding.

By modifying the method of Proposition 3.3 we can now show that the change of metric converges in C^0 . For $t \in (-\infty, 0]$, define $f_t(y_0) = e^{i\beta t} \cdot g_t(y_0)$ and let $t_1 \leq t_2 \leq T < 0$. This is summarised in the diagram below.

Proposition 3.17. $h_t(y_0)$ converges in the C^0 norm to a unique limit $h_\infty(y_0) \in \mathcal{G}^{\mathbb{C}}/\mathcal{G}$ as $t \rightarrow -\infty$. The limit depends continuously on the initial condition $y_0 \in W_x^-$. The rate of convergence is given by

$$(3.15) \quad \sup_X \sigma(h_t(y_0)(h_\infty(y_0))^{-1}) \leq C_2 e^{2\eta t} \|y_0 - x\|_{L_k^2}$$



where $C_2 > 0$ is a constant depending only on the orbit $\mathcal{G} \cdot x$, the constant η is from Proposition 2.21 and k is a positive integer chosen so that $L_k^2 \hookrightarrow C^1$ is a continuous embedding.

Proof. The result follows from the same procedure as the proof of Proposition 3.3, except now we use the estimate from Lemma 3.16 instead of the estimate from Lemma 3.2 and the distance-decreasing formula for the modified flow from Lemma 2.19.

Let $t_1 < t_2 \leq T < 0$, and let $h_{t_1-t_2}(y_{t_2})$ be the change of metric connecting $y_{t_2} = g_{t_2}(y_0) \cdot y_0$ and $e^{i\beta(t_1-t_2)} \cdot y_{t_1}$. Lemma 3.16 and the estimate (3.14) above show that $h_{t_1-t_2}(y_{t_2})$ satisfies

$$\begin{aligned} \sup_X \sigma(h_{t_1-t_2}(y_{t_2})) &\leq C \left(\|e^{i\beta(t_1-t_2)} \cdot y_{t_1} - x\|_{C^1} + \|y_{t_2} - x\|_{C^1} \right) \\ &\leq CC' \|y_{t_2} - x\|_{L_k^2} + C \|y_{t_2} - x\|_{C^1} \\ &\leq C'' \|y_T - x\|_{L_k^2} \leq C_2 e^{2\eta T} \|y_0 - x\|_{L_k^2} \end{aligned}$$

By the construction of the modified flow, the gauge transformation connecting y_{t_2} and $e^{i\beta(t_1-t_2)} \cdot y_{t_1}$ is in \mathcal{G}_*^C . The distance-decreasing formula for the action of $e^{i\beta(t_1-t_2)}$ from Lemma 2.19 then implies that


$$\sigma(h_{t_1}(y_0)h_{t_2}(y_0)^{-1}) \leq \sigma(h_{t_1-t_2}(y_{t_2}))$$

and so the sequence $h_t(y_0)$ is Cauchy in the C^0 norm, by the same proof as Proposition 3.3. \square

Therefore y_0 is connected to z_∞^0 by a C^0 gauge transformation. Elliptic regularity together with the fact that z_∞^0 is a C^∞ Higgs pair then shows that y_0 is gauge equivalent to z_∞^0 by a C^∞ gauge transformation.

The same method as the proof of Proposition 3.14 then allows us to explicitly construct a solution of the linearised flow $z_\infty^{-s} = e^{i\beta s} \cdot z_\infty^0$ converging to x as $s \rightarrow +\infty$. More precisely, for $t < 0$ define $y_t = g_t(y_0) \cdot y_0$ and $z_t^0 = e^{i\beta t} \cdot y_t$ (recall that $t < 0$ is used for the upwards flow). Then (3.13) shows that there exists a subsequence $\{t_n\}$ such that $z_{t_n}^0 \rightarrow z_\infty^0$ strongly in L_{k-1}^2 . Given any $t < -s < 0$, define $y_{-s} = g_{-s}(y_0) \cdot y_0$ and $z_{t+s}^{-s} := e^{i\beta(t+s)} \cdot g_{t+s}(y_{-s}) \cdot y_{-s} = e^{i\beta(t+s)} \cdot y_t$, and note that $z_{t_n}^0 = e^{-i\beta s} \cdot z_{t_n+s}^{-s}$.

Then for each s there exists a further subsequence $\{t_{n_\ell}\}$ such that $z_{t_{n_\ell}+s}^{-s} \rightarrow z_\infty^{-s}$ strongly in L_{k-1}^2 . Since $z_{t_{n_\ell}+s}^{-s} = e^{i\beta s} \cdot z_{t_{n_\ell}}^0$, $z_{t_{n_\ell}}^0 \rightarrow z_\infty^0$ and $z_{t_{n_\ell}+s}^{-s} \rightarrow z_\infty^{-s}$ then the continuity of the action of $e^{i\beta s}$ implies that



$z_\infty^{-s} = e^{i\beta s} \cdot z_\infty^0$. Replacing y_0 by y_{-s} in the estimate (3.13) shows that $z_\infty^{-s} \rightarrow x$ as $s \rightarrow \infty$. Lemma 2.10 then shows that z_∞^0 is $\mathcal{G}^{\mathbb{C}}$ equivalent to a point in S_x^- , which is smooth by Lemma 2.7.

Therefore y_0 is $\mathcal{G}^{\mathbb{C}}$ equivalent to a point in S_x^- , and so we have proved the converse to Proposition 3.14.

Proposition 3.18. *For each $y_0 \in W_x^-$ there exists a C^∞ gauge transformation $g \in \mathcal{G}^{\mathbb{C}}$ such that $g \cdot y_0 \in S_x^-$.*

3.3. An algebraic criterion for the existence of flow lines. The results of the previous two sections combine to give the following theorem.

Theorem 3.19. *Let E be a complex vector bundle over a compact Riemann surface X , and let $(\bar{\partial}_A, \phi)$ be a Higgs bundle on E . Suppose that E admits a filtration $(E^{(1)}, \phi^{(1)}) \subset \dots \subset (E^{(n)}, \phi^{(n)}) = (E, \phi)$ by Higgs subbundles such that the quotients $(Q_k, \phi_k) := (E^{(k)}, \phi^{(k)}) / (E^{(k-1)}, \phi^{(k-1)})$ are Higgs polystable and $\text{slope}(Q_k) < \text{slope}(Q_j)$ for all $k < j$. Then there exists $g \in \mathcal{G}^{\mathbb{C}}$ and a solution to the reverse Yang-Mills-Higgs heat flow equation with initial condition $g \cdot (\bar{\partial}_A, \phi)$ which converges to a critical point isomorphic to $(Q_1, \phi_1) \oplus \dots \oplus (Q_n, \phi_n)$.*

Conversely, if there exists a solution of the reverse heat flow from the initial condition $(\bar{\partial}_A, \phi)$ converging to a critical point $(Q_1, \phi_1) \oplus \dots \oplus (Q_n, \phi_n)$ then $(\bar{\partial}_A, \phi)$ admits a filtration $(E^{(1)}, \phi^{(1)}) \subset \dots \subset (E^{(n)}, \phi^{(n)}) = (E, \phi)$ whose graded object is isomorphic to $(Q_1, \phi_1) \oplus \dots \oplus (Q_n, \phi_n)$.

Proof. Suppose first that $(\bar{\partial}_A, \phi)$ admits a filtration $(E^{(1)}, \phi^{(1)}) \subset \dots \subset (E^{(n)}, \phi^{(n)}) = (E, \phi)$ by Higgs subbundles such that the quotients $(Q_k, \phi_k) := (E^{(k)}, \phi^{(k)}) / (E^{(k-1)}, \phi^{(k-1)})$ are Higgs polystable and $\text{slope}(Q_k) < \text{slope}(Q_j)$ for all $k < j$. Let x be a critical point isomorphic to $(Q_1, \phi_1) \oplus \dots \oplus (Q_n, \phi_n)$, and let U be the neighbourhood of x from Lemma 2.10. Then by applying the isomorphism $x \cong (Q_1, \phi_1) \oplus \dots \oplus (Q_n, \phi_n)$ and scaling the extension classes there exists a complex gauge transformation such that $g \cdot (\bar{\partial}_A, \phi)$ is in U . Applying Lemma 2.10 shows that $(\bar{\partial}_A, \phi)$ is isomorphic to a point in S_x^- , and therefore Proposition 3.14 shows that $(\bar{\partial}_A, \phi)$ is isomorphic to a point in W_x^- .

Conversely, if $x = (Q_1, \phi_1) \oplus \cdots \oplus (Q_n, \phi_n)$ is a critical point and $(\bar{\partial}_A, \phi) \in W_x^-$, then Proposition 3.18 shows that there exists $g \in \mathcal{G}^{\mathbb{C}}$ such that $g \cdot (\bar{\partial}_A, \phi) \in S_x^-$. Therefore Lemma 2.10 shows that $(\bar{\partial}_A, \phi)$ admits a filtration whose graded object is isomorphic to $(Q_1, \phi_1) \oplus \cdots \oplus (Q_n, \phi_n)$. \square

4. THE HECKE CORRESPONDENCE VIA YANG-MILLS-HIGGS FLOW LINES

Let (E, ϕ) be a polystable Higgs bundle of rank r and degree d , and let (L_u, ϕ_u) be a Higgs line bundle with $\deg L_u < \text{slope } E$. Let F be a smooth complex vector bundle C^∞ isomorphic to $E \oplus L_u$ and choose a metric on F such that the Higgs structure on $(E, \phi) \oplus (L_u, \phi_u)$ is a Yang-Mills-Higgs critical point in the space $\mathcal{B}(F)$ of Higgs bundles on F . The goal of this section is to show that Hecke modifications of the Higgs bundle (E, ϕ) correspond to Yang-Mills-Higgs flow lines in $\mathcal{B}(F)$ connecting the critical point $(E, \phi) \oplus (L_u, \phi_u)$ to lower critical points.

In Section 4.1 we review Hecke modifications of Higgs bundles. Section 4.2 describes how the space of Hecke modifications relates to the geometry of the negative slice and Section 4.3 contains the proof of Theorem 4.22 which shows that Hecke modifications correspond to YMH flow lines. In Section 4.4 we give a geometric criterion for points to be connected by unbroken flow lines in terms of the secant varieties of the space of Hecke modifications inside the negative slice. In particular, this gives a complete classification of the YMH flow lines for rank 2 (cf. Corollary 4.27). Throughout this section the notation \mathcal{E} is used to denote the sheaf of holomorphic sections of the bundle E .

4.1. Hecke modifications of Higgs bundles. The purpose of this section is to derive some basic results for Hecke modifications of Higgs bundles which will be used in Section 4.3 to prove Theorem 4.22. In Section 4.4 we extend these results to study unbroken YMH flow lines. The Hecke correspondence was introduced by Narasimhan and Ramanan [36], who applied it to study automorphisms of the moduli space of stable bundles [37] and desingularisations of this space [38]. One important aspect of the Hecke correspondence is its role in the Geometric Langlands program (cf. [10] and [31]; an excellent survey is [14]). It also appears in the physics approach to Geometric Langlands [25] which has been further explained in [52].

First recall that a Hecke modification of a holomorphic bundle E over a Riemann surface X is determined by points $p_1, \dots, p_n \in X$ (not necessarily distinct) and nonzero elements $v_j \in E_{p_j}^*$ for $j = 1, \dots, n$. This data determines a sheaf homomorphism $\mathcal{E} \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j}$ to the skyscraper sheaf supported at p_1, \dots, p_n with kernel a locally free sheaf \mathcal{E}' . This determines a holomorphic bundle $E' \rightarrow X$ which we call the *Hecke modification of E determined by $v = (v_1, \dots, v_n)$* .

$$0 \rightarrow \mathcal{E}' \rightarrow \mathcal{E} \xrightarrow{v} \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$$

Since the kernel sheaf \mathcal{E}' only depends on the equivalence class of each v_j in $\mathbb{P}E_{p_j}^*$ then from now on we abuse the notation slightly and also use $v_j \in \mathbb{P}E_{p_j}^*$ to denote the equivalence class of $v_j \in E_{p_j}^*$.

As explained in [52, Sec. 4.5], if (E, ϕ) is a Higgs bundle, then a Hecke modification of (E, ϕ) may introduce poles into the Higgs field and so there are restrictions on the allowable modifications which preserve holomorphicity of the Higgs field.

Definition 4.1. Let (E, ϕ) be a Higgs bundle. A Hecke modification E' of E is *compatible* with ϕ if the induced Higgs field on E' is holomorphic.

The next result describes a basic condition for the modification to be compatible with the Higgs field.

Lemma 4.2. *Let (E, ϕ) be a Higgs bundle, and $0 \rightarrow \mathcal{E}' \rightarrow \mathcal{E} \xrightarrow{v} \mathbb{C}_p \rightarrow 0$ a Hecke modification of E induced by $v \in E_p^*$. Then the induced Higgs field ϕ' on E' is holomorphic if and only if there exists an eigenvalue μ of $\phi(p)$ such that the composition $\mathcal{E} \otimes K^{-1} \xrightarrow{\phi - \mu \cdot \text{id}} \mathcal{E} \xrightarrow{v} \mathbb{C}_p$ is zero.*

Proof. Let $\phi \in H^0(\text{End}(E) \otimes K)$. Then ϕ pulls back to a holomorphic Higgs field $\phi' \in H^0(\text{End}(E') \otimes K)$ if and only if for any open set $U \subset X$ and any section $s \in \mathcal{E}(U)$, the condition $s \in \ker(\mathcal{E}(U) \xrightarrow{v} \mathbb{C}_p(U))$ implies that $\phi(s) \in \ker((\mathcal{E} \otimes K)(U) \xrightarrow{v} \mathbb{C}_p(U))$. After choosing a trivialisation of K in a neighbourhood of p , we can decompose the Higgs field $\phi(p)$ on the fibre E_p as follows

$$(4.1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \ker v & \longrightarrow & E_p & \longrightarrow & \mathbb{C}_p \longrightarrow 0 \\ & & \downarrow \phi(p)|_{\ker v} & & \downarrow \phi(p) & & \downarrow \mu \\ 0 & \longrightarrow & \ker v & \longrightarrow & E_p & \longrightarrow & \mathbb{C}_p \longrightarrow 0 \end{array}$$

where scalar multiplication by μ is induced from the action of $\phi(p)$ on the quotient $\mathbb{C}_p = E_p / \ker v$. Therefore the endomorphism $(\phi(p) - \mu \cdot \text{id})$ maps E_p into the subspace $\ker v$ and so $v \in E_p^*$ descends to a well-defined homomorphism $v' : \text{coker}(\phi(p) - \mu \cdot \text{id}) \rightarrow \mathbb{C}$.

Conversely, given an eigenvalue μ of $\phi(p)$ and an element $v' \in \text{coker}(\phi(p) - \mu \cdot \text{id})^*$, one can choose a basis of E_p and extend v' to an element $v \in E_p^*$ such that $\text{im}(\phi(p) - \mu \cdot \text{id}) \subset \ker v$. Equivalently, $\phi(p)$ preserves $\ker v$ and so $v \in E_p^*$ defines a Hecke modification E' of E such that the induced Higgs field on E' is holomorphic. \square

Corollary 4.3. *Let (E, ϕ) be a Higgs bundle and let $0 \rightarrow \mathcal{E}' \rightarrow \mathcal{E} \xrightarrow{v} \mathbb{C}_p \rightarrow 0$ be a Hecke modification of E induced by $v \in \mathbb{P}E_p^*$. The following conditions are equivalent*

- (1) *The induced Higgs field ϕ' on E' is holomorphic.*
- (2) *There exists an eigenvalue μ of $\phi(p)$ such that $v(\phi(s)) = \mu v(s)$ for all sections s of E .*
- (3) *There exists an eigenvalue μ of $\phi(p)$ such that v descends to a well-defined $v' \in (\text{coker}(\phi(p) - \mu \cdot \text{id}))^*$.*

Lemma 4.4. *Let (E, ϕ) be a Higgs bundle and (G, φ) a Higgs subsheaf. Then there exists a Higgs subbundle $(G', \varphi') \subset (E, \phi)$ such that $\text{rank}(G) = \text{rank}(G')$ and (G, φ) is a Higgs subsheaf of (G', φ') .*

Proof. Since $\dim_{\mathbb{C}} X = 1$ then a standard procedure shows that there is a holomorphic subbundle $G' \subset E$ with $\text{rank}(G) = \text{rank}(G')$ and G is a subsheaf of G' , and so it only remains to show that this is a Higgs subbundle. The reverse of the construction above shows that the Higgs field φ preserving G extends to a meromorphic Higgs field φ' preserving G' , and since this is the restriction of a holomorphic Higgs field ϕ on E to the holomorphic subbundle G' , then φ' must be holomorphic on G' . Therefore G' is ϕ -invariant. \square

Definition 4.5. A Higgs bundle (E, ϕ) is (m, n) -stable (resp. (m, n) -semistable) if for every proper ϕ -invariant holomorphic subbundle $F \subset E$ we have

$$\frac{\deg F + m}{\text{rank } F} < \frac{\deg E - n}{\text{rank } E} \quad (\text{resp. } \leq).$$

If (E, ϕ) is $(0, n)$ -semistable then any Hecke modification $0 \rightarrow (\mathcal{E}', \phi') \rightarrow (\mathcal{E}, \phi) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$ is semistable.

Definition 4.6. Then *space of admissible Hecke modifications* is the subset $\mathcal{N}_\phi \subset \mathbb{P}E^*$ corresponding to the Hecke modifications which are compatible with the Higgs field.

Remark 4.7. (1) If $\phi = 0$ then $\mathcal{N}_0 = \mathbb{P}E^*$. If E is $(0, 1)$ -stable then there is a well-defined map $\mathbb{P}E^* \rightarrow \mathbb{P}H^1(E^*)$. The construction of the next section generalises this to a map $\mathcal{N}_\phi \rightarrow \mathbb{P}\mathcal{H}^1(E^*)$ (cf. Remark 4.16).

(2) Note that the construction above is the reverse of that described in [52], which begins with E' and modifies the bundle to produce a bundle E with $\deg E = \deg E' + 1$. Here we begin with E and construct E' via a modification $0 \rightarrow \mathcal{E}' \rightarrow \mathcal{E} \rightarrow \mathbb{C}_p \rightarrow 0$ since we want to interpret the compatible modifications in terms of the geometry of the negative slice (see Section 4.2) in order to draw a connection with the results on gradient flow lines for the Yang-Mills-Higgs flow functional from Section 3.3.

(3) One can also see the above construction more explicitly in local coordinates as in [52] by choosing a local frame $\{s_1, \dots, s_n\}$ for E in a neighbourhood U of p with local coordinate z centred at p and for which the evaluation map $\mathcal{E} \xrightarrow{v} \mathbb{C}_p$ satisfies $v(s_1) = s_1(0)$ and $v(s_j) = 0$ for all $j = 2, \dots, n$. Then over $U \setminus \{p\}$, the functions $\{\frac{1}{z}s_1(z), s_2(z), \dots, s_n(z)\}$ form a local frame for E' . Equivalently, the transition function $g = \begin{pmatrix} \frac{1}{z} & 0 \\ 0 & \text{id} \end{pmatrix}$ maps the trivialisation for E to a trivialisation for E' (note that this is the inverse of the transition function from [52, Sec. 4.5.2] for the reason explained in the previous paragraph). In this local frame field on E we write $\phi(z) = \begin{pmatrix} A(z) & B(z) \\ C(z) & D(z) \end{pmatrix}$. The action on the Higgs field is then

$$g \begin{pmatrix} A(z) & B(z) \\ C(z) & D(z) \end{pmatrix} g^{-1} = \begin{pmatrix} A(z) & \frac{1}{z}B(z) \\ zC(z) & D(z) \end{pmatrix}$$

Therefore the induced Higgs field on E' will have a pole at p unless $B(0) = 0$. The scalar $A(0)$ in this local picture is the same as the scalar μ from (4.1), and we see that

$$\phi(p) - \mu \cdot \text{id} = \begin{pmatrix} 0 & 0 \\ C(0) & D(0) - \mu \cdot \text{id} \end{pmatrix}$$

With respect to the basis of E_p given by the choice of local frame, $v(\phi(p) - \mu \cdot \text{id}) = 0$. Moreover, via this local frame $\text{coker}(\phi(p) - \mu \cdot \text{id})$ is identified with a subspace of E_p which contains the linear span of $s_1(0)$. Therefore we see in the local coordinate picture that $v \in E_p^*$ descends to an element of $(\text{coker}(\phi(p) - \mu \cdot \text{id}))^*$.

The next result shows that the admissible Hecke modifications have an interpretation in terms of the spectral curve associated to the Higgs field. This extends the results of [52] to include the possibility that p is a branch point of the spectral cover.

First recall Hitchin's construction of the spectral curve from [16]. Let (E, ϕ) be a Higgs pair. Then there is a projection map $\pi : K \rightarrow X$ and a bundle π^*E over the total space of the canonical bundle together with a tautological section λ of π^*E . The zero set of the characteristic polynomial of $\pi^*\phi$ defines a subvariety S inside the total space of K . The projection π restricts to a map $\pi : S \rightarrow X$, where for each $p \in X$ the fibre $\pi^{-1}(p)$ consists of the eigenvalues of the Higgs field $\phi(p)$. As explained in [16], generically the discriminant of the Higgs field has simple zeros and in this case S is a smooth curve called the *spectral curve*. The induced projection $\pi : S \rightarrow X$ is then a ramified covering map with ramification divisor denoted $\mathcal{R} \subset S$.

The pullback of the Higgs field to the spectral curve is a bundle homomorphism $\pi^*E \rightarrow \pi^*(E \otimes K)$, and the eigenspaces correspond to $\ker(\pi^*\phi - \lambda \cdot \text{id})$, where λ is the tautological section defined above. When the discriminant of the Higgs field has simple zeros then Hitchin shows in [16] that the eigenspaces form a line bundle $\mathcal{N} \rightarrow S$ and that the original bundle E can be reconstructed as $\pi_*\mathcal{L}$, where the line bundle $\mathcal{L} \rightarrow S$ is formed by modifying \mathcal{N} at the ramification points $0 \rightarrow \mathcal{N} \rightarrow \mathcal{L} \rightarrow \bigoplus_{p \in \mathcal{R}} \mathbb{C}_p \rightarrow 0$. One can reconstruct the Higgs field ϕ by pushing forward the endomorphism defined by the tautological section $\lambda : \mathcal{L} \rightarrow \mathcal{L} \otimes \pi^*K$.

Lemma 4.8. *If the discriminant of ϕ has simple zeros then an admissible Hecke modification of (E, ϕ) corresponds to a Hecke modification of the line bundle \mathcal{L} over the spectral curve.*

Proof. Consider the pullback bundle $\pi^*E \rightarrow S$. The pullback of the Higgs field induces a sheaf homomorphism $(\pi^*\phi - \lambda \cdot \text{id}) : \pi^*\mathcal{E} \otimes (\pi^*K)^{-1} \rightarrow \pi^*\mathcal{E}$. As explained in [52, Sec. 2.6], when the discriminant of ϕ has simple zeros then the cokernel of this homomorphism is the line bundle $\mathcal{L} \rightarrow S$ such that $\mathcal{E} \cong \pi_*\mathcal{L}$.

For $\mu \in S$ such that $p = \pi(\mu)$, there is an isomorphism of the stalks of the skyscraper sheaves $\mathbb{C}_p \cong \pi_*(\mathbb{C}_\mu)$. Then a Hecke modification $\mathcal{L} \xrightarrow{v'} \mathbb{C}_\mu$ given by nonzero $v' \in \mathcal{L}_\mu^*$ induces a Hecke modification $v = v' \circ q \circ \pi^* : \mathcal{E} \rightarrow \mathbb{C}_p$, defined by the commutative diagram below.

$$\begin{array}{ccccccc}
 \pi^*\mathcal{E} \otimes (\pi^*K)^{-1} & \xrightarrow{\pi^*\phi - \lambda \cdot \text{id}} & \pi^*\mathcal{E} & \xrightarrow{q} & \text{coker}(\pi^*\phi - \lambda \cdot \text{id}) & \longrightarrow & 0 \\
 & & \uparrow \pi^* & & \searrow v' & & \\
 & & \mathcal{E} & \xrightarrow{v} & \mathbb{C}_p & \longrightarrow & 0
 \end{array}$$

The definition of v implies that for any open set $U \subset X$ with a trivialisation of K in a neighbourhood of p , and all $s \in \mathcal{E}(U)$ we have

$$v(\phi s) = v' \circ q(\pi^*(\phi s)) = v' \circ q(\mu \pi^*(s)) = \mu v' \circ q \circ \pi^*(s) = \mu v(s)$$

and so v is compatible with the Higgs field by Corollary 4.3.

Conversely, let $v \in E_p^*$ be compatible with the Higgs field ϕ . Corollary 4.3 shows that this induces a well-defined element of $\text{coker}(\phi - \mu \cdot \text{id})^*$. Consider the endomorphisms $\phi(p) - \mu \cdot \text{id}$ on the fibre of E over $p \in X$ and $\pi^*\phi(\mu) - \mu \cdot \text{id}$ on the fibre of π^*E over $\mu \in S$.

$$\begin{array}{ccccccc}
(\pi^* E \otimes \pi^* K^{-1})_\mu & \xrightarrow{\pi^* \phi - \mu \cdot \text{id}} & (\pi^* E)_\mu & \longrightarrow & \text{coker}(\pi^* \phi - \mu \cdot \text{id})_\mu & \longrightarrow & 0 \\
\uparrow & & \uparrow & & \uparrow & & \\
(E \otimes K^{-1})_p & \xrightarrow{\phi - \mu \cdot \text{id}} & E_p & \longrightarrow & \text{coker}(\phi - \mu \cdot \text{id})_p & \longrightarrow & 0
\end{array}$$

The universal property of cokernel defines a map $\text{coker}(\phi - \mu \cdot \text{id})_p \rightarrow \text{coker}(\pi^* \phi - \mu \cdot \text{id})_\mu$. Since the discriminant of the Higgs field has simple zeros then both fibres are one-dimensional and so this map becomes an isomorphism. Therefore v induces a well-defined homomorphism on the fibre $\text{coker}(\pi^* \phi - \mu \cdot \text{id})_\mu \rightarrow \mathbb{C}$, and hence a Hecke modification of \mathcal{L} at $\mu \in S$. \square

Remark 4.9. When $p \in X$ is not a branch point of $\pi : S \rightarrow X$ then this result is contained in [52].

Corollary 4.10. *If the discriminant of ϕ has simple zeros then the space of Hecke modifications is $\mathcal{N}_\phi = S$.*

4.2. Secant varieties associated to the space of Hecke modifications. The purpose of this section is to connect the geometry of the space of Hecke modifications with the geometry of the negative slice at a critical point in order to prepare for the proof of Theorem 4.22 in the next section.

Let (E_1, ϕ_1) and (E_2, ϕ_2) be Higgs bundles and let $\bar{\partial}_A$ denote the induced holomorphic structure on $E_1^* E_2$. Then there is an elliptic complex

$$\Omega^0(E_1^* E_2) \xrightarrow{L_1} \Omega^{0,1}(E_1^* E_2) \oplus \Omega^{1,0}(E_1^* E_2) \xrightarrow{L_2} \Omega^{1,1}(E_1^* E_2),$$

where $L_1(u) = (\bar{\partial}_A u, \phi_2 u - u \phi_1)$ and $L_2(a, \varphi) = (\bar{\partial}_A \varphi + [a, \phi])$. Let $\mathcal{H}^0 = \ker L_1$, $\mathcal{H}^1 = \ker L_1^* \cap \ker L_2$ and $\mathcal{H}^2 = \ker L_2^*$ denote the spaces of harmonic forms. Recall that if (E_1, ϕ_1) and (E_2, ϕ_2) are both Higgs stable and $\text{slope}(E_2) < \text{slope}(E_1)$ then $\mathcal{H}^0(E_1^* E_2) = 0$.

Now consider the special case where (E_1, ϕ_1) is $(0, n)$ -stable and (E_2, ϕ_2) is a Higgs line bundle. Let \mathcal{B} denote the space of Higgs bundles on the smooth bundle $E_1 \oplus E_2$ and choose a metric such that $(E_1, \phi_1) \oplus (E_2, \phi_2)$ is a critical point of $\text{YMH} : \mathcal{B} \rightarrow \mathbb{R}$. Definition 2.5 shows that $\mathcal{H}^1(E_1^* E_2) \cong S_x^-$ is the negative slice at this critical point.

Let $0 \rightarrow (\mathcal{E}', \phi') \rightarrow (\mathcal{E}_1, \phi_1) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$ be a Hecke modification defined by $v_1, \dots, v_n \in \mathbb{P}E_1^*$. Applying the functor $\text{Hom}(\cdot, \mathcal{E}_2)$ to the short exact sequence $0 \rightarrow \mathcal{E}' \rightarrow \mathcal{E}_1 \rightarrow \bigoplus_j \mathbb{C}_{p_j} \rightarrow 0$ gives us an exact sequence of sheaves $0 \rightarrow \text{Hom}(\mathcal{E}_1, \mathcal{E}_2) \rightarrow \text{Hom}(\mathcal{E}', \mathcal{E}_2) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j}^* \rightarrow 0$, where the final term comes from the isomorphism $\text{Ext}^1(\bigoplus_j \mathbb{C}_{p_j}, \mathcal{E}_2) \cong \text{Hom}(\mathcal{E}_2, \bigoplus_j \mathbb{C}_{p_j} \otimes K)^* \cong \bigoplus_j \mathbb{C}_{p_j}^*$. Note that this depends on a choice of trivialisations of E_2 and K , however the kernel of the map $\text{Hom}(\mathcal{E}', \mathcal{E}_2) \rightarrow \bigoplus_j \mathbb{C}_{p_j}$ is independent of these choices. This gives us the following short exact sequence of Higgs sheaves

$$(4.2) \quad 0 \rightarrow \mathcal{E}_1^* \mathcal{E}_2 \rightarrow (\mathcal{E}')^* \mathcal{E}_2 \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j}^* \rightarrow 0$$

There is an induced map $\Omega^0((\mathcal{E}')^* \mathcal{E}_2) \rightarrow \Omega^{1,0}((\mathcal{E}')^* \mathcal{E}_2)$ given by $s \mapsto \phi_2 s - s \phi'$. Recall from Corollary 4.3 that there exists an eigenvalue μ_j for $\phi_1(p_j)$ such that $v(\phi_1(p_j) - \mu_j \cdot \text{id}) = 0$ for each $j = 1, \dots, n$. From the above exact sequence there is an induced homomorphism $\Omega^{1,0}((\mathcal{E}')^* \mathcal{E}_2) \xrightarrow{ev^1} \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$. The component of $ev^1(\phi_2 s - s \phi')$ in \mathbb{C}_{p_j} is $(\phi_2(p_j) - \mu_j)s$. In particular, $\phi_2 s - s \phi' \in \ker(ev^1)$ iff $\phi_2(p_j) = \mu_j$ for all $j = 1, \dots, n$.

Definition 4.11. Let (E_1, ϕ_1) be a Higgs bundle, and (E_2, ϕ_2) a Higgs line bundle. The *space of Hecke modifications compatible with ϕ_1 and ϕ_2* , denoted $\mathcal{N}_{\phi_1, \phi_2} \subset \mathcal{N}_{\phi_1}$, is the set of Hecke modifications compatible with ϕ_1 such that $ev^1(\phi_2 s - s\phi') = 0$ for all $s \in \Omega^0((E')^*E_2)$.

Remark 4.12. Note that if $n = 1$ and $v \in \mathbb{P}E_1^*$ is a Hecke modification compatible with ϕ_1 , then the requirement that $v \in \mathcal{N}_{\phi_1, \phi_2}$ reduces to $\phi_2(p) = \mu$, where μ is the eigenvalue of $\phi_1(p)$ from Corollary 4.3. Such a $\phi_2 \in H^0(\text{End}(E_2) \otimes K) = H^0(K)$ always exists since the canonical linear system is basepoint free and therefore $\bigcup_{\phi_2 \in H^0(K)} \mathcal{N}_{\phi_1, \phi_2} = \mathcal{N}_{\phi_1}$. If $n > 1$ then ϕ_2 with these properties may not exist for some choices of $\phi_1 \in H^0(\text{End}(E_1) \otimes K)$ and $v_1, \dots, v_n \in \mathbb{P}E_1^*$ (the existence of ϕ_2 depends on the complex structure of the surface X). If $\phi_1 = 0$, then we can choose $\phi_2 = 0$ and in this case $\mathcal{N}_{\phi_1, \phi_2} = \mathcal{N}_{\phi_1} = \mathbb{P}E_1^*$ (this corresponds to the case of the Yang-Mills flow in Theorem 4.22).

Lemma 4.13. Let (E_1, ϕ_1) be Higgs polystable and (E_2, ϕ_2) be a Higgs line bundle. Let $0 \rightarrow (\mathcal{E}', \phi') \rightarrow (\mathcal{E}, \phi) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$ be a Hecke modification defined by distinct $v_1, \dots, v_n \in \mathcal{N}_{\phi_1, \phi_2}$.

Then there is an exact sequence

$$(4.3) \quad 0 \rightarrow \mathcal{H}^0(E_1^*E_2) \rightarrow \mathcal{H}^0((E')^*E_2) \rightarrow \mathbb{C}^n \rightarrow \mathcal{H}^1(E_1^*E_2) \rightarrow \mathcal{H}^1((E')^*E_2)$$

Proof. The short exact sequence (4.2) leads to the following commutative diagram of spaces of smooth sections

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega^0(E_1^*E_2) & \xrightarrow{i^*} & \Omega^0((E')^*E_2) & \xrightarrow{ev^0} & \bigoplus_{j=1}^n \mathbb{C}_{p_j} \longrightarrow 0 \\ & & \downarrow L_1 & & \downarrow L_1 & & \\ 0 & \longrightarrow & \Omega^{0,1}(E_1^*E_2) \oplus \Omega^{1,0}(E_1^*E_2) & \xrightarrow{i^*} & \Omega^{0,1}((E')^*E_2) \oplus \Omega^{1,0}((E')^*E_2) & \xrightarrow{ev^1} & \bigoplus_{j=1}^n \mathbb{C}_{p_j} \oplus \mathbb{C}_{p_j} \longrightarrow 0 \end{array}$$

Since $\bar{\partial}_A s$ depends on the germ of a section around a point, then there is no well-defined map $\bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \oplus \mathbb{C}_{p_j}$ making the diagram commute, so the exact sequence (4.3) does not follow immediately from the standard construction, and therefore we give an explicit construction below.

First construct a map $\mathbb{C}^n \rightarrow \mathcal{H}^1(E_1^*E_2)$ as follows. Given $z \in \mathbb{C}^n$, choose a smooth section $s' \in \Omega^0((E')^*E_2)$ such that $ev^0(s') = z$ and $ev^1(\bar{\partial}_A s') = 0$. Since $\phi_2(p_j) = \mu_j$, then $ev^1(\phi_2 s' - s'\phi') = 0$ and so $ev^1(L_1 s') = 0$. Therefore $(\bar{\partial}_A s', \phi_2 s' - s'\phi') = i^*(a, \varphi)$ for some $(a, \varphi) \in \Omega^{0,1}(E_1^*E_2) \oplus \Omega^{1,0}(E_1^*E_2)$. Let $[(a, \varphi)] \in \mathcal{H}^1(E_1^*E_2)$ denote the harmonic representative of (a, φ) . Define the map $\mathbb{C}^n \rightarrow \mathcal{H}^1(E_1^*E_2)$ by $z \mapsto [(a, \varphi)]$.

To see that this is well-defined independent of the choice of $s' \in \Omega^0((E')^*E_2)$, note that if $s'' \in \Omega^0((E')^*E_2)$ is another section such that $ev^0(s'') = z$ and $ev^1(\bar{\partial}_A s'') = 0$, then $ev^0(s'' - s') = 0$, and so $s'' - s' = i^*(s)$ for some $s \in \Omega^0(E_1^*E_2)$. Therefore $L_1(s'' - s') = i^*L_1(s)$ with $[L_1(s)] = 0 \in \mathcal{H}^1(E_1^*E_2)$, and so s' and s'' determine the same harmonic representative in $\mathcal{H}^1(E_1^*E_2)$.

To check exactness of (4.3) at the term \mathbb{C}^n , note that if $z = ev^0(s')$ for some harmonic $s' \in \mathcal{H}^0((E')^*E_2)$, then $L_1(s') = 0 = i^*(0, 0)$, and so $z \in \mathbb{C}^n$ maps to $0 \in \mathcal{H}^1(E_1^*E_2)$. Moreover, if z maps to $0 \in \mathcal{H}^1(E_1^*E_2)$, then there exists $s' \in \Omega^0((E')^*E_2)$ such that $L_1(s') = i^*(a, \varphi)$ where $(a, \varphi) \in \Omega^{0,1}(E_1^*E_2) \oplus \Omega^{1,0}((E')^*E_2)$ and $(a, \varphi) = L_1(s)$ for some $s \in \Omega^0(E_1^*E_2)$. Therefore s' and i^*s differ by a harmonic

section of $\mathcal{H}^0((E')^*E_2)$. Since $ev^0(i^*s) = 0$ then z is the image of this harmonic section under the map $\mathcal{H}^0((E')^*E_2) \rightarrow \mathbb{C}^n$.

To check exactness at $\mathcal{H}^1(E_1^*E_2)$, given $z \in \mathbb{C}^n$ construct (a, φ) as above and note that $i^*(a, \varphi) = L_1s'$ for some $s' \in \Omega^0((E')^*E_2)$. Therefore $i^*[(a, \varphi)] = 0 \in \mathcal{H}^1((E')^*E_2)$ and so the image of $\mathbb{C}^n \rightarrow \mathcal{H}^1(E_1^*E_2)$ is contained in the kernel of $\mathcal{H}^1(E_1^*E_2) \rightarrow \mathcal{H}^1((E')^*E_2)$. Now suppose that the image of $[(a, \varphi)]$ is zero in $\mathcal{H}^1((E')^*E_2)$, i.e. $i^*(a, \varphi) = L_1s'$ for some $s' \in \Omega^0((E')^*E_2)$. Let $z = ev^0(s')$. Note that $z = 0$ implies that $s' = i^*s$ for some $s \in \Omega^0(E_1^*E_2)$, and so $[(a, \varphi)] = 0$. If $z \neq 0$ then there exists $s'' \in \Omega^0((E')^*E_2)$ such that $ev^1(L_1(s'')) = 0$ and $ev^0(s'') = z$. Then $L_1(s'') = i^*(a'', \varphi'')$ for some $(a'', \varphi'') \in \Omega^{0,1}(E_1^*E_2) \oplus \Omega^{1,0}(E_1^*E_2)$. Moreover, $ev^0(s'' - s') = 0$, so $s'' - s' = i^*s$ for some $s \in \Omega^0(E_1^*E_2)$. Commutativity implies that $L_1s = (a'', \varphi'') - (a, \varphi)$, and so the harmonic representatives $[(a, \varphi)]$ and $[(a'', \varphi'')]$ are equal. Therefore $[(a, \varphi)]$ is the image of z by the map $\mathbb{C}^n \rightarrow \mathcal{H}^1(E_1^*E_2)$, which completes the proof of exactness at $\mathcal{H}^1(E_1^*E_2)$.

Exactness at the rest of the terms in the sequence (4.3) then follows from standard methods. \square

For any stable Higgs bundle (E, ϕ) with $d = \deg E$ and $r = \text{rank } E$, define the *generalised Segre invariant* by

$$s_k(E, \phi) := kd - r \left(\max_{F \subset E, \text{rank } F=k} \deg F \right).$$

where the maximum is taken over all ϕ -invariant holomorphic subbundles of rank k . Note that $s_k(E, \phi) \geq s_k(E, 0) =: s_k(E)$ and

$$\frac{1}{rk} s_k(E, \phi) = \min_{F \subset E, \text{rank } F=k} (\text{slope}(E) - \text{slope}(F))$$

Note that any Hecke modification $(E', \phi') \hookrightarrow (E, \phi)$ with $\deg E - \deg E' = n$ has Segre invariant $s_k(E', \phi') \geq s_k(E, \phi) - nk$. As a special case, (E', ϕ') is stable if $n < \frac{1}{k}s_k(E, \phi)$ for all $k = 1, \dots, r-1$.

A theorem of Lange [30, Satz 2.2] shows that a general stable holomorphic bundle E satisfies $s_k(E) \geq k(r-k)(g-1)$ for all $k = 1, \dots, r-1$. Since there is a dense open subset of stable Higgs bundles whose underlying holomorphic bundle is stable, then Lange's theorem also gives the same lower bound on the Segre invariant for a general stable Higgs bundle.

Lemma 4.14. *Let $0 \rightarrow (E', \phi') \rightarrow (E, \phi) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$ be a Hecke modification defined by distinct points $v_1, \dots, v_n \in \mathbb{P}E^*$ such that $n < \frac{1}{k}s_k(E, \phi)$ for all $k = 1, \dots, r-1$. Then $\text{slope}(G) < \text{slope}(E')$ for any proper non-zero Higgs subbundle $(G, \phi_G) \subset (E, \phi)$. In particular, this condition is satisfied if (E, ϕ) is a general stable Higgs bundle and $n < g-1$.*

Proof. Let $k = \text{rank } G$ and $h = \deg G$. Then the lower bound on the Segre invariant implies that

$$\begin{aligned} \text{slope}(E') - \text{slope}(G) &= \frac{d-n}{r} - \frac{h}{k} = \frac{1}{rk} (kd - kn - rh) \\ &\geq \frac{1}{rk} (s_k(E, \phi) - kn) \end{aligned}$$

Therefore if $n < \frac{1}{k}s_k(E, \phi)$ then $\text{slope}(E') - \text{slope}(G) > 0$ for any Higgs subbundle of rank k . If $n < g-1$ then [30, Satz 2.2] shows that this condition is satisfied for general stable Higgs bundles. \square

Corollary 4.15. *Let (E_1, ϕ_1) be a stable Higgs bundle, let $n < \frac{1}{k}s_k(E_1, \phi_1)$ for all $k = 1, \dots, \text{rank}(E_1) - 1$ and let (E_2, ϕ_2) be a Higgs line bundle such that $\deg E_2 < \frac{\deg E_1 - n}{\text{rank } E_1}$. Then given any set of n distinct points $\{v_1, \dots, v_n\} \subset \mathcal{N}_{\phi_1, \phi_2}$ there is a well-defined n -dimensional subspace $\ker(\mathcal{H}^1(E_1^*E_2) \rightarrow \mathcal{H}^1((E')^*E_2))$.*

Proof. Let (E', ϕ') be the Hecke modification of (E_1, ϕ_1) determined by $\{v_1, \dots, v_n\} \subset \mathbb{P}E_1^*$. The lower bound on the Segre invariant implies that (E', ϕ') is Higgs stable, and therefore $\mathcal{H}^0((E')^*E_2) = 0$ since $\text{slope}(E_2) < \text{slope}(E') = \frac{\deg E_1 - n}{\text{rank } E_1}$. The exact sequence (4.3) then reduces to

$$0 \rightarrow \mathbb{C}^n \rightarrow \mathcal{H}^1(E_1^*E_2) \rightarrow \mathcal{H}^1((E')^*E_2)$$

and so $\ker(\mathcal{H}^1(E_1^*E_2) \rightarrow \mathcal{H}^1((E')^*E_2))$ is a well-defined n -dimensional subspace of $\mathcal{H}^1(E_1^*E_2)$ associated to $\{v_1, \dots, v_n\}$. \square

Remark 4.16. As noted above, the maps $\mathbb{C}^n \rightarrow \mathcal{H}^1(E_1^*E_2)$ depend on choosing trivialisations, but different choices lead to the same map up to a change of basis of \mathbb{C}^n , and so the subspace $\ker(\mathcal{H}^1((E')^*E_2) \rightarrow \mathcal{H}^1(E_1^*E_2))$ is independent of these choices.

In the special case where $n = 1$, then this construction gives a well-defined map $\mathcal{N}_{\phi_1, \phi_2} \rightarrow \mathbb{P}\mathcal{H}^1(E_1^*E_2)$. When $n < \frac{1}{k}s_k(E_1, \phi_1)$ for all k , then Corollary 4.15 shows that any n distinct points v_1, \dots, v_n span a nondegenerate copy of \mathbb{P}^{n-1} in $\mathbb{P}\mathcal{H}^1(E_1^*E_2)$.

In the special case where $\phi_1 = \phi_2 = 0$ and E_2 is trivial, then $\mathcal{N}_{\phi_1, \phi_2} = \mathbb{P}E^*$ and $\mathcal{H}^1(E_1^*) \cong H^{0,1}(E_1^*) \oplus H^{1,0}(E_1^*)$. Then the map $\mathbb{P}E^* \rightarrow \mathcal{H}^1(E_1^*) \rightarrow H^{0,1}(E_1^*) \cong H^0(E_1 \otimes K)^*$ is the usual map defined for holomorphic bundles (cf. [21, p804]).

Definition 4.17. The n^{th} secant variety, denoted $\text{Sec}^n(\mathcal{N}_{\phi_1, \phi_2}) \subset \mathbb{P}\mathcal{H}^1(E_1^*E_2)$, is the union of the subspaces $\text{span}\{v_1, \dots, v_n\} \subset \mathbb{P}\mathcal{H}^1(E_1^*E_2)$ taken over all n -tuples of distinct points $v_1, \dots, v_n \in \mathcal{N}_{\phi_1, \phi_2}$.

The next lemma is a Higgs bundle version of [35, Lemma 3.1]. Since the proof is similar to that in [35] then it is omitted.

Lemma 4.18. *Let $0 \rightarrow (E_2, \phi_2) \rightarrow (F, \tilde{\phi}) \rightarrow (E_1, \phi_1) \rightarrow 0$ be an extension of Higgs bundles defined by the extension class $[(a, \varphi)] \in \mathcal{H}^1(E_1^*E_2)$. Let $(E', \phi') \xrightarrow{i} (E_1, \phi_1)$ be a Higgs subsheaf such that $i^*[(a, \varphi)] = 0 \in \mathcal{H}^1((E')^*E_2)$. Then (E', ϕ') is a Higgs subsheaf of $(F, \tilde{\phi})$.*

$$\begin{array}{ccccccc}
 & & & & (E', \phi') & & \\
 & & & & \swarrow & \downarrow i & \\
 0 & \longrightarrow & (E_2, \phi_2) & \longrightarrow & (F, \tilde{\phi}) & \longrightarrow & (E_1, \phi_1) \longrightarrow 0
 \end{array}$$

Corollary 4.19. *Let (E_1, ϕ_1) be stable, let $n < \frac{1}{k}s_k(E_1, \phi_1)$ for all $k = 1, \dots, \text{rank}(E_1) - 1$, let (E_2, ϕ_2) be a Higgs line bundle and suppose that $\deg E_2 < \frac{\deg E_1 - n}{\text{rank } E_1}$. Let $0 \rightarrow (E_2, \phi_2) \rightarrow (F, \tilde{\phi}) \rightarrow (E_1, \phi_1) \rightarrow 0$ be an extension of Higgs bundles with extension class $[(a, \varphi)] \in \mathcal{H}^1(E_1^*E_2)$. Let $0 \rightarrow (E', \phi') \xrightarrow{i} (E_1, \phi_1) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$ be a Hecke modification determined by distinct points $\{v_1, \dots, v_n\} \in \mathcal{N}_{\phi_1, \phi_2}$.*

*Then (E', ϕ') is a subsheaf of $(F, \tilde{\phi})$ if $[(a, \varphi)] \in \text{span}\{v_1, \dots, v_n\} \subset \mathcal{H}^1(E_1^*E_2)$.*

Proof. If $[(a, \varphi)] \in \text{span}\{v_1, \dots, v_n\}$ then $[(a, \varphi)] \in \ker(\mathcal{H}^1(E_1^*E_2) \rightarrow \mathcal{H}^1((E')^*E_2))$ by Corollary 4.15, and therefore (E', ϕ') is a subsheaf of $(F, \tilde{\phi})$ by Lemma 4.18. \square

The next lemma gives a condition on the extension class $[(a, \varphi)] \in \mathcal{H}^1(E_1^*E_2)$ for (E', ϕ') to be the subsheaf of largest degree which lifts to a subsheaf of $(F, \tilde{\phi})$. This is used to study unbroken flow lines in Section 4.4.

Lemma 4.20. *Let (E_1, ϕ_1) be a stable Higgs bundle, choose n such that $2n - 1 < \frac{1}{k}s_k(E_1, \phi_1)$ for all $k = 1, \dots, \text{rank}(E_1)$, let (E_2, ϕ_2) be a Higgs line bundle and suppose that $\deg E_2 < \frac{\deg E_1 - (2n-1)}{\text{rank } E_1}$. Let $0 \rightarrow (E_2, \phi_2) \rightarrow (F, \tilde{\phi}) \rightarrow (E_1, \phi_1) \rightarrow 0$ be an extension of Higgs bundles with extension class $[(a, \varphi)] \in \text{Sec}^n(\mathcal{N}_{\phi_1, \phi_2}) \setminus \text{Sec}^{n-1}(\mathcal{N}_{\phi_1, \phi_2}) \subset \mathbb{P}\mathcal{H}^1(E_1^*E_2)$ and let $0 \rightarrow (E', \phi') \xrightarrow{i} (E_1, \phi_1) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$ be a Hecke modification determined by distinct points $v_1, \dots, v_n \in \mathcal{N}_{\phi_1, \phi_2}$ such that $i^*[(a, \varphi)] = 0$.*

Let $(\mathcal{E}'', \phi'') \xrightarrow{i''} (\mathcal{E}, \phi)$ be a subsheaf such that $(i'')^[(a, \varphi)] = 0 \in \mathcal{H}^1((E'')^*E_2)$ and $\text{rank } E'' = \text{rank } E$. Then $\deg(E'') \leq \deg(E')$.*

Proof. Let $\{v'_1, \dots, v'_m\} \subset \mathcal{N}_{\phi_1, \phi_2}$ be the set of distinct points defining the Hecke modification $(\mathcal{E}'', \phi'') \xrightarrow{i''} (E_1, \phi_1)$. Then $i^*[(a, \varphi)] = 0$ and $(i'')^*[(a, \varphi)] = 0$ together imply that $[(a, \varphi)] \in \text{span}\{v_1, \dots, v_n\} \cap \text{span}\{v'_1, \dots, v'_m\}$. Either $m + n > 2n - 1$ (and so $\deg E'' \leq \deg E'$) or $m + n \leq 2n - 1$ in which case Corollary 4.15 together with the lower bound $2n - 1 < \frac{1}{k}s_k(E_1, \phi_1)$ implies that $\text{span}\{v_1, \dots, v_n\} \cap \text{span}\{v'_1, \dots, v'_m\}$ is the linear span of $\{v_1, \dots, v_n\} \cap \{v'_1, \dots, v'_m\}$. Since $m + n \leq 2n - 1$ then $\{v_1, \dots, v_n\} \cap \{v'_1, \dots, v'_m\}$ is a strict subset of $\{v_1, \dots, v_n\}$, which is not possible since $[(a, \varphi)] \notin \text{Sec}^{n-1}(\mathcal{N}_{\phi_1, \phi_2})$. Therefore $\deg E'' \leq \deg E'$. \square

4.3. Constructing Hecke modifications of Higgs bundles via the Yang-Mills-Higgs flow. Let (E, ϕ) be a stable Higgs bundle and L_u a line bundle with $\deg L_u < \frac{\deg E - 1}{\text{rank } E}$, and let E' be a Hecke modification of E which is compatible with the Higgs field

$$0 \rightarrow (E', \phi') \xrightarrow{i} (\mathcal{E}, \phi) \xrightarrow{v} \mathbb{C}_p \rightarrow 0.$$

The goal of this section is to construct critical points $x_u = (L_u, \phi_u) \oplus (E, \phi)$ and $x_\ell = (L_\ell, \phi_\ell) \oplus (E', \phi')$ together with a broken flow line connecting x_u and x_ℓ . The result of Theorem 3.19 shows that this amounts to constructing a Higgs field $\phi_u \in H^0(K)$, a Higgs pair $(F, \tilde{\phi})$ in the unstable set of x_u and a complex gauge transformation $g \in \mathcal{G}^{\mathbb{C}}$ such that (E', ϕ') is a Higgs subbundle of $g \cdot (F, \tilde{\phi})$.

Lemma 4.21. *Let $0 \rightarrow (E', \phi') \rightarrow (\mathcal{E}, \phi) \xrightarrow{v} \mathbb{C}_p \rightarrow 0$ be a Hecke modification such that (E, ϕ) and (E', ϕ') are both Higgs semistable, and let L_u be a line bundle with $\deg L_u < \text{slope}(E') < \text{slope}(E)$. Then there exists a Higgs field $\phi_u \in H^0(K)$ and a non-trivial Higgs extension $(F, \tilde{\phi})$ of (L_u, ϕ_u) by (E, ϕ) such that (E', ϕ') is a Higgs subsheaf of $(F, \tilde{\phi})$.*

Proof. By Remark 4.12, there exists $\phi_u \in H^0(K)$ such that $v \in \mathcal{N}_{\phi, \phi_u}$. Since (E', ϕ') is semistable with $\text{slope}(E') > \text{slope}(L_u)$ then $\mathcal{H}^0((E')^*L_u) = 0$ and so the exact sequence (4.3) shows that the Hecke modification $v \in \mathbb{P}E^*$ determines a one-dimensional subspace of $\mathcal{H}^1(E^*L_u)$, and that any non-trivial extension class in this subspace is in the kernel of the map $\mathcal{H}^1(E^*L_u) \rightarrow \mathcal{H}^1((E')^*L_u)$. Let

$0 \rightarrow (L_u, \phi_u) \rightarrow (F, \tilde{\phi}) \rightarrow (E, \phi) \rightarrow 0$ be such an extension. Then Lemma 4.18 shows that (E', ϕ') is a Higgs subsheaf of $(F, \tilde{\phi})$. \square

We can now use this result to relate Hecke modifications at a single point with YMH flow lines.

- Theorem 4.22.** (1) *Let $0 \rightarrow (E', \phi') \rightarrow (E, \phi) \xrightarrow{v} \mathbb{C}_p \rightarrow 0$ be a Hecke modification such that (E, ϕ) is stable and (E', ϕ') is semistable, and let L_u be a line bundle with $\deg L_u + 1 < \text{slope}(E') < \text{slope}(E)$. Then there exist sections $\phi_u, \phi_\ell \in H^0(K)$, a line bundle L_ℓ with $\deg L_\ell = \deg L_u + 1$ and a metric on $E \oplus L_u$ such that $x_u = (E, \phi) \oplus (L_u, \phi_u)$ and $x_\ell = (E'_{gr}, \phi'_{gr}) \oplus (L_\ell, \phi_\ell)$ are critical points connected by a YMH flow line, where (E'_{gr}, ϕ'_{gr}) is isomorphic to the graded object of the Seshadri filtration of (E', ϕ') .*
- (2) *Let $x_u = (E, \phi) \oplus (L_u, \phi_u)$ and $x_\ell = (E', \phi') \oplus (L_\ell, \phi_\ell)$ be critical points connected by a YMH flow line such that L_u, L_ℓ are line bundles with $\deg L_u = \deg L_\ell + 1$, (E, ϕ) is stable and (E', ϕ') is polystable with $\deg L_u + 1 < \text{slope}(E') < \text{slope}(E)$. If (E', ϕ') is Higgs stable then it is a Hecke modification of (E, ϕ) . If (E', ϕ') is Higgs polystable then it is the graded object of the Seshadri filtration of a Hecke modification of (E, ϕ) .*

Proof of Theorem 4.22. Given a Hecke modification $0 \rightarrow (E', \phi') \rightarrow (E, \phi) \rightarrow \mathbb{C}_p \rightarrow 0$ as in Lemma 4.21, choose $\phi_u \in H^0(K)$ such that $v \in \mathcal{N}_{\phi, \phi_u}$ and apply a gauge transformation to $E \oplus L_u$ such that $x_u = (E, \phi) \oplus (L_u, \phi_u)$ is a critical point of YMH. The harmonic representative of the extension class $[(a, \varphi)] \in \mathcal{H}^1(E^* L_u)$ from Lemma 4.21 defines an extension $0 \rightarrow (L_u, \phi_u) \rightarrow (F, \tilde{\phi}) \rightarrow (E, \phi) \rightarrow 0$ such that $y = (F, \tilde{\phi})$ is in the negative slice of x_u , and therefore flows down to a limit isomorphic to the graded object of the Harder-Narasimhan-Seshadri filtration of $(F, \tilde{\phi})$.


Lemma 4.21 also shows that (E', ϕ') is a Higgs subsheaf of $(F, \tilde{\phi})$. Lemma 4.4 shows that this has a resolution as a Higgs subbundle of $(F, \tilde{\phi})$, however since the Harder-Narasimhan type of $(F, \tilde{\phi})$ is strictly less than that of $(E, \phi) \oplus (L_u, \phi_u)$, $\text{rank}(E') = \text{rank}(F) - 1$ and $\deg E' = \deg E - 1$, then (E', ϕ') already has the maximal possible slope for a semistable Higgs subbundle of $(F, \tilde{\phi})$, and therefore (E', ϕ') must be the maximal semistable Higgs subbundle. Since $\text{rank}(E') = \text{rank}(F) - 1$, then the graded object of the Harder-Narasimhan-Seshadri filtration of $(F, \tilde{\phi})$ is $(E'_{gr}, \phi'_{gr}) \oplus (L_\ell, \phi_\ell)$, where $(L_\ell, \phi_\ell) = (F, \tilde{\phi}) / (E', \phi')$. Theorem 3.19 then shows that $(E, \phi) \oplus (L_u, \phi_u)$ and $(E'_{gr}, \phi'_{gr}) \oplus (L_\ell, \phi_\ell)$ are connected by a flow line.

Conversely, if $x_u = (E, \phi) \oplus (L_u, \phi_u)$ and $x_\ell = (E', \phi') \oplus (L_\ell, \phi_\ell)$ are critical points connected by a flow line, then Theorem 3.19 shows that there exists a Higgs pair $(F, \tilde{\phi})$ in the negative slice of x_u such that (E', ϕ') is the graded object of the Seshadri filtration of the maximal semistable Higgs subbundle of $(F, \tilde{\phi})$. If (E', ϕ') is Higgs stable, then since $\text{slope}(E') > \text{slope}(L_u)$ we see (E', ϕ') is a Higgs subsheaf of (E, ϕ) with $\text{rank}(E) = \text{rank}(E')$ and $\deg(E') = \deg(E) - 1$. Therefore (E', ϕ') is a Hecke modification of (E, ϕ) . If (E', ϕ') is Higgs polystable then the same argument shows that (E', ϕ') is the graded object of the Seshadri filtration of a Hecke modification of (E, ϕ) . \square

In general, for any flow one can define the space $\mathcal{F}_{\ell, u}$ of flow lines connecting upper and lower critical sets C_u and C_ℓ , and the space $\mathcal{P}_{\ell, u} \subset C_u \times C_\ell$ of pairs of critical points connected by a flow line. These spaces are equipped with projection maps to the critical sets defined by the canonical projection taking a


flow line to its endpoints.

(4.4)



For the Yang-Mills-Higgs flow, given critical sets C_u and C_ℓ of respective Harder-Narasimhan types $(\frac{d}{r}, \deg L_u)$ and $(\frac{d-1}{r}, \deg L_u + 1)$ as in Theorem 4.22 above, there are natural projection maps to the moduli space $C_u \rightarrow \mathcal{M}_{ss}^{Higgs}(r, d)$ and $C_\ell \rightarrow \mathcal{M}_{ss}^{Higgs}(r, d-1)$. Since the flow is \mathcal{G} -equivariant, then there is an induced correspondence variety $\mathcal{M}_{\ell,u} \subset \mathcal{M}_{ss}^{Higgs}(r, d-1) \times \mathcal{M}_{ss}^{Higgs}(r, d)$.

(4.5)



Theorem 4.22 shows that $((E', \phi'), (E, \phi)) \in \mathcal{M}_{\ell,u}$ if and only if (E', ϕ') is a Hecke modification of (E, ϕ) and both Higgs pairs are semistable. If r and d are coprime then $\mathcal{M}_{ss}^{Higgs}(r, d)$ consists of stable Higgs pairs and so every Hecke modification of (E, ϕ) is semistable. Therefore we have proved

Corollary 4.23. $\mathcal{M}_{\ell,u}$ is the Hecke correspondence.

For Hecke modifications defined at multiple points (non-miniscule Hecke modifications in the terminology of [52]), we immediately have the following result.

Corollary 4.24. *Let (E, ϕ) be a $(0, n)$ -stable Higgs bundle and consider a Hecke modification $0 \rightarrow (\mathcal{E}', \phi') \rightarrow (\mathcal{E}, \phi) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_n} \rightarrow 0$ defined by $n > 1$ distinct points $\{v_1, \dots, v_n\} \in \mathbb{P}E^*$. If there exists $\phi_u \in H^0(K)$ such that $v_1, \dots, v_n \in \mathcal{N}_{\phi, \phi_u}$, then there is a broken flow line connecting $x_u = (E, \phi) \oplus (L_u, \phi_u)$ and $x_\ell = (E'_{gr}, \phi'_{gr}) \oplus (L_\ell, \phi_\ell)$, where (E'_{gr}, ϕ'_{gr}) is the graded object of the Seshadri filtration of the semistable Higgs bundle (E', ϕ') .*

Proof. Inductively apply Theorem 4.22. □

4.4. A geometric criterion for unbroken YMH flow lines. Corollary 4.24 gives a criterion for two YMH critical points $x_u = (E, \phi) \oplus (L_u, \phi_u)$ and $x_\ell = (E', \phi') \oplus (L_\ell, \phi_\ell)$ to be connected by a broken flow line. It is natural to ask whether they are also connected by an *unbroken* flow line. The goal of this section is to answer this question by giving a geometric construction for points in the negative slice of x_u which correspond to unbroken flow lines connecting x_u and x_ℓ in terms of the secant varieties $\text{Sec}^n(\mathcal{N}_{\phi, \phi_u})$. For holomorphic bundles, the connection between secant varieties and Hecke modifications has been studied in [29], [4] and [17].

Given a YMH critical point $x_u = (E, \phi) \oplus (L_u, \phi_u)$ with (E, ϕ) stable and $\text{rank } L_u = 1$, consider an extension $0 \rightarrow (L_u, \phi_u) \rightarrow (F, \tilde{\phi}) \rightarrow (E, \phi) \rightarrow 0$ with extension class $[(a, \varphi)] \in \mathcal{H}^1(E^* L_u) = S_{x_u}^-$. Let $0 \rightarrow (E', \phi') \rightarrow (E, \phi) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$ be a Hecke modification of (E, ϕ) as in the previous lemma, such that $\text{deg } L_u < \text{slope}(E')$.

Lemma 4.25. *If (G, ϕ_G) is a semistable Higgs subbundle of $(F, \tilde{\phi})$ with $\text{slope}(G) > \text{deg } L_u$ and $\text{rank}(G) < \text{rank}(E)$, then there is a Higgs subbundle $(G', \phi'_G) \subset (E, \phi)$ with $\text{slope}(G') \geq \text{slope}(G)$ and $\text{rank}(G) = \text{rank}(G')$.*

Proof. If (G, ϕ_G) is a semistable Higgs subbundle of $(F, \tilde{\phi})$ with $\text{slope}(G) > \text{deg } L_u$, then $\mathcal{H}^0(G^* L_u) = 0$, and so (G, ϕ_G) is a Higgs subsheaf of (E, ϕ) .

$$\begin{array}{ccccccc}
 & & & & (G, \phi_G) & & \\
 & & & & \swarrow & \downarrow & \\
 0 & \longrightarrow & (L_u, \phi_u) & \longrightarrow & (F, \tilde{\phi}) & \longrightarrow & (E, \phi) \longrightarrow 0
 \end{array}$$

Lemma 4.4 shows that the subsheaf (G, ϕ_G) can be resolved to form a Higgs subbundle (G', ϕ'_G) of (E, ϕ) with $\text{slope}(G') \geq \text{slope}(G)$. \square

Theorem 4.26. *Let (E, ϕ) be a stable Higgs bundle with Segre invariant $s_k(E, \phi)$ and choose n such that $0 < 2n - 1 < \min_{1 \leq k \leq r-1} (\frac{1}{k} s_k(E, \phi))$. Let $0 \rightarrow (\mathcal{E}', \phi') \rightarrow (\mathcal{E}, \phi) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0$ be a Hecke modification of (E, ϕ) defined by distinct points $v_1, \dots, v_n \in \mathbb{P}E^*$, and let (L_u, ϕ_u) be a Higgs line bundle such that $v_1, \dots, v_n \in \mathcal{N}_{\phi, \phi_u}$. Choose a metric such that $x_u = (E, \phi) \oplus (L_u, \phi_u)$ is a YMH critical point.*

Then any extension class $[(a, \varphi)] \in \text{span}\{v_1, \dots, v_n\} \cap (\text{Sec}^n(\mathcal{N}_{\phi, \phi_u}) \setminus \text{Sec}^{n-1}(\mathcal{N}_{\phi, \phi_u})) \subset \mathbb{P}\mathcal{H}^1(E^ L_u)$ is isomorphic to an unbroken flow line connecting $x_u = (E, \phi) \oplus (L_u, \phi_u)$ and $x_\ell = (E', \phi') \oplus (L_\ell, \phi_\ell)$.*

Proof. Let $(F, \tilde{\phi})$ be a Higgs bundle determined by the extension class $[(a, \varphi)] \in \mathbb{P}\mathcal{H}^1(E^* L_u)$. The choice of bundle is not unique, but the isomorphism class of $(F, \tilde{\phi})$ is unique. The proof reduces to showing that (E', ϕ') is the maximal semistable Higgs subbundle of $(F, \tilde{\phi})$.

Since $[(a, \varphi)] \notin \text{Sec}^{n-1}(\mathcal{N}_{\phi, \phi_u})$, then Lemma 4.20 shows that (E', ϕ') is the subsheaf of (E, ϕ) with maximal degree among those that lift to a subsheaf of $(F, \tilde{\phi})$. Any semistable Higgs subbundle (E'', ϕ'') of $(F, \tilde{\phi})$ with $\text{rank}(E'') = \text{rank}(E)$ either has $\text{slope}(E'') \leq \text{deg } L_u < \text{slope}(E')$, or it is a subsheaf of (E, ϕ) and so must have $\text{slope}(E'') \leq \text{slope}(E')$.

The previous lemma shows that if (G, ϕ_G) is any semistable Higgs subbundle of $(F, \tilde{\phi})$ with $\text{slope}(G) > \text{deg } L_u$ and $\text{rank}(G) < \text{rank}(E)$, then there is a Higgs subbundle (G', ϕ'_G) of (E, ϕ) with $\text{slope}(G') \geq \text{slope}(G)$. The upper bound on $n = \text{deg } E - \text{deg } E'$ in terms of the Segre invariant then implies that $\text{slope}(E') > \text{slope}(G') \geq \text{slope}(G)$ by Lemma 4.14.

Therefore the subbundle $(\tilde{E}', \tilde{\phi}')$ resolving the subsheaf $(E', \phi') \subset (F, \tilde{\phi})$ is the maximal semistable Higgs subbundle of $(F, \tilde{\phi})$. Since $(\tilde{E}', \tilde{\phi}')$ is semistable and $\text{slope}(\tilde{E}') \geq \text{slope}(E') > \text{deg } L_u$, then $\mathcal{H}^0((\tilde{E}')^* L_u) = 0$, and so $(\tilde{E}', \tilde{\phi}')$ is a Higgs subsheaf of (E, ϕ) that lifts to a subbundle of $(F, \tilde{\phi})$. Since $\text{deg } E'$ is maximal among all such subsheaves, then we must have $(E', \phi') = (\tilde{E}', \tilde{\phi}')$ and so (E', ϕ') is the maximal semistable subbundle of $(F, \tilde{\phi})$. Therefore Theorem 3.19 shows that x_u and x_ℓ are connected by an unbroken flow line. \square

If $\text{rank}(F) = 2$ (so that E is a line bundle), then the condition on the Segre invariant $s_k(E, \phi)$ becomes vacuous. Moreover, $\mathbb{P}E^* \cong X$ and so Hecke modifications of E are determined by a subset $\{v_1, \dots, v_n\} \subset X$. Therefore in the case $\text{rank}(F) = 2$, we have a complete classification of the YMH flow lines on the space of Higgs bundles $\mathcal{B}(F)$.

Corollary 4.27. *Let $F \rightarrow X$ be a C^∞ Hermitian vector bundle with $\text{rank}(F) = 2$. Let $x_u = (L_1^u, \phi_1^u) \oplus (L_2^u, \phi_2^u)$ and $x_\ell = (L_1^\ell, \phi_1^\ell) \oplus (L_2^\ell, \phi_2^\ell)$ be non-minimal critical points with $\text{YMH}(x_u) > \text{YMH}(x_\ell)$. Suppose without loss of generality that $\deg L_1^u > \deg L_1^\ell > \deg L_2^\ell > \deg L_2^u$. Let $n = \deg L_1^u - \deg L_1^\ell$.*

Then x_u and x_ℓ are connected by a broken flow line if and only if there exists $\{v_1, \dots, v_n\} \in \mathcal{N}_{\phi_1^u, \phi_2^u}$ such that

$$\begin{aligned} 0 &\rightarrow (L_1^\ell, \phi_1^\ell) \rightarrow (L_1^u, \phi_1^u) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0 \\ 0 &\rightarrow (L_2^u, \phi_2^u) \rightarrow (L_2^\ell, \phi_2^\ell) \rightarrow \bigoplus_{j=1}^n \mathbb{C}_{p_j} \rightarrow 0 \end{aligned}$$

are both Hecke modifications determined by $\{v_1, \dots, v_n\}$. They are connected by an unbroken flow line if the previous condition holds and $\{v_1, \dots, v_n\} \in \text{Sec}^n(\mathcal{N}_{\phi_1^u, \phi_2^u}) \setminus \text{Sec}^{n-1}(\mathcal{N}_{\phi_1^u, \phi_2^u})$.

APPENDIX A. UNIQUENESS FOR THE REVERSE YANG-MILLS-HIGGS FLOW

The methods of Donaldson [8] and Simpson [45] show that the Yang-Mills-Higgs flow resembles a non-linear heat equation, and therefore the backwards flow is ill-posed. In Section 3.1 we prove existence of solutions to the backwards heat flow that converge to a critical point. To show that these solutions are well-defined we prove in this section that if a solution to the reverse YMH flow exists then it must be unique.

Using the Hermitian metric, let d_A be the Chern connection associated to $\bar{\partial}_A$ and let $\psi = \phi + \phi^* \in \Omega^1(i \text{ad}(E))$. The holomorphicity condition $\bar{\partial}_A \phi = 0$ becomes the pair of equations $d_A \psi = 0$, $d_A^* \psi = 0$ which also imply that $[F_A, \psi] = d_A^2 \psi = 0$, and the Yang-Mills-Higgs functional is $\|F_A + \psi \wedge \psi\|_{L^2}^2$.

Proposition A.1. *Let $(d_{A_1}, \psi_1)(t)$, $(d_{A_2}, \psi_2)(t)$ be two solutions of the Yang-Mills-Higgs flow (A.4) on a compact Riemann surface with respective initial conditions $(d_{A_1}, \psi_1)(0)$ and $(d_{A_2}, \psi_2)(0)$. If there exists a finite $T > 0$ such that $(d_{A_1}, \psi_1)(T) = (d_{A_2}, \psi_2)(T)$ then $(d_{A_1}, \psi_1)(t) = (d_{A_2}, \psi_2)(t)$ for all $t \in [0, T]$.*

The result of Proposition A.1 is valid when the base manifold is a compact Riemann surface, since we use the estimates of [51, Sec. 3.2] to prove that the constant C in Lemma A.2 is uniform. In the case of the Yang-Mills flow on a compact Kähler manifold the estimates of Donaldson in [8] show that we can make this constant uniform on a finite time interval $[0, T]$ and so the result also applies in this setting. The setup described in the previous paragraph consisting of Higgs pairs (d_A, ψ) satisfying $d_A \psi = 0$, $d_A^* \psi = 0$ is valid on any Riemannian manifold, and so the result of Proposition A.1 will also apply to any class of solutions for which one can prove that the connection, Higgs field, the curvature and all of their derivatives are uniformly bounded on the given finite time interval $[0, T]$.

Let ∇_A denote the covariant derivative associated to the connection d_A . The complex connection associated to the pair (d_A, ψ) is $D_{(A, \psi)} \eta = d_A \eta + [\psi, \eta]$ and the Laplacian is $\Delta_{(A, \psi)} \eta = D_{(A, \psi)}^* D_{(A, \psi)} \eta + D_{(A, \psi)} D_{(A, \psi)}^* \eta$ for any form $\eta \in \Omega^p(\text{End}(E))$. The equation $d_A \psi = 0$ implies that the curvature of the complex connection is $D_{(A, \psi)} D_{(A, \psi)} \eta = [F_A + \psi \wedge \psi, \eta]$.

We have the following identities which will be useful in what follows. The notation $a \times b$ is used to denote various bilinear expressions with constant coefficients.

$$(A.1) \quad 0 = d_A(F_A + \psi \wedge \psi), \quad 0 = [\psi, F_A + \psi \wedge \psi]$$

$$(A.2) \quad \Delta_{(A,\psi)}\eta = \nabla_A^* \nabla_A \eta + (F_A + \psi \wedge \psi) \times \eta + R_M \times \eta + \psi \times \psi \times \eta + \nabla_A \psi \times \psi \times \eta$$

$$(A.3) \quad 0 = D_{(A,\psi)}^* D_{(A,\psi)}^*(F_A + \psi \wedge \psi)$$

The first identity follows from the Bianchi identity and the equation $d_A \psi = 0$. Equation (A.2) is the Weitzenböck identity for a Higgs pair which follows from the usual identity for ∇_A (see for example [3]) together with the fact that $(\psi \wedge \psi) \times \eta$ and the remaining terms in the Laplacian are of the form $\psi \times \psi \times \eta + \nabla_A \psi \times \psi \times \eta$. To see the identity (A.3), take the inner product of the right hand side with an arbitrary $\eta \in \Omega^0(\text{End}(E))$. We have (cf. [41, (2.2)] for the case $\psi = 0$)

$$\begin{aligned} \left\langle D_{(A,\psi)}^* D_{(A,\psi)}^*(F_A + \psi \wedge \psi), \eta \right\rangle &= \left\langle F_A + \psi \wedge \psi, D_{(A,\psi)} D_{(A,\psi)} \eta \right\rangle \\ &= \left\langle F_A + \psi \wedge \psi, [F_A + \psi \wedge \psi, \eta] \right\rangle = 0 \end{aligned}$$

Consider the Yang-Mills-Higgs flow equations

$$(A.4) \quad \frac{\partial A}{\partial t} = -d_A^*(F_A + \psi \wedge \psi), \quad \frac{\partial \psi}{\partial t} = *[\psi, *(F_A + \psi \wedge \psi)]$$

After using the metric to decompose $\Omega^1(\text{End}(E)) \cong \Omega^1(\text{ad}(E)) \oplus \Omega^1(i \text{ad}(E))$, the flow equation can be written more compactly as

$$\frac{\partial}{\partial t}(d_A + \psi) = -D_{(A,\psi)}^*(F_A + \psi \wedge \psi)$$

We then have

$$\begin{aligned} \frac{\partial}{\partial t}(F_A + \psi \wedge \psi) &= d_A \left(\frac{\partial A}{\partial t} \right) + \frac{\partial \psi}{\partial t} \wedge \psi + \psi \wedge \frac{\partial \psi}{\partial t} \\ &= -d_A d_A^*(F_A + \psi \wedge \psi) + [\psi, *[\psi, *(F_A + \psi \wedge \psi)]] \\ &= -\Delta_{(A,\psi)}(F_A + \psi \wedge \psi) - d_A *[\psi, *(F_A + \psi \wedge \psi)] + [\psi, d_A^*(F_A + \psi \wedge \psi)] \end{aligned}$$

where in the last step we use the Bianchi identity (A.1). We also have

$$\begin{aligned} \frac{\partial}{\partial t}(d_A^*(F_A + \psi \wedge \psi)) &= - * \left[\frac{\partial A}{\partial t}, *(F_A + \psi \wedge \psi) \right] + d_A^* \left(\frac{\partial}{\partial t}(F_A + \psi \wedge \psi) \right) \\ &= * [d_A^*(F_A + \psi \wedge \psi), F_A + \psi \wedge \psi] - d_A^* d_A d_A^*(F_A + \psi \wedge \psi) \\ &\quad + d_A^* [\psi, *[\psi, *(F_A + \psi \wedge \psi)]] \end{aligned}$$

and

$$\begin{aligned} \frac{\partial}{\partial t}(- * [\psi, *(F_A + \psi \wedge \psi)]) &= - * \left[\frac{\partial \psi}{\partial t}, *(F_A + \psi \wedge \psi) \right] - * \left[\psi, \frac{\partial}{\partial t} *(F_A + \psi \wedge \psi) \right] \\ &= * [- * [\psi, *(F_A + \psi \wedge \psi)], *(F_A + \psi \wedge \psi)] + * [\psi, * d_A d_A^*(F_A + \psi \wedge \psi)] \\ &\quad - * [\psi, *[\psi, *[\psi, *(F_A + \psi \wedge \psi)]]] \end{aligned}$$

Adding these two results gives us

$$\begin{aligned} \frac{\partial}{\partial t} \left(D_{(A,\psi)}^*(F_A + \psi \wedge \psi) \right) &= * \left[D_{(A,\psi)}^*(F_A + \psi \wedge \psi), F_A + \psi \wedge \psi \right] - D_{(A,\psi)}^* D_{(A,\psi)} D_{(A,\psi)}^*(F_A + \psi \wedge \psi) \\ &= * \left[D_{(A,\psi)}^*(F_A + \psi \wedge \psi), F_A + \psi \wedge \psi \right] - \Delta_{(A,\psi)} D_{(A,\psi)}^*(F_A + \psi \wedge \psi) \end{aligned}$$

where the last step uses (A.3). Let $\mu_{(A,\psi)} = F_A + \psi \wedge \psi$ and $\nu_{(A,\psi)} = D_{(A,\psi)}^*(F_A + \psi \wedge \psi)$. The above equations become

$$(A.5) \quad \left(\frac{\partial}{\partial t} + \Delta_{(A,\psi)} \right) \mu_{(A,\psi)} = -d_A * [\psi, *(F_A + \psi \wedge \psi)] + [\psi, d_A^*(F_A + \psi \wedge \psi)]$$

$$(A.6) \quad \left(\frac{\partial}{\partial t} + \Delta_{(A,\psi)} \right) \nu_{(A,\psi)} = *[\nu_{(A,\psi)}, *\mu_{(A,\psi)}]$$

Now consider two solutions $(A_1, \psi_1)(t)$ and $(A_2, \psi_2)(t)$ to the Yang-Mills-Higgs flow equations (A.4) on the time interval $[0, T]$ such that $(A_1, \psi_1)(T) = (A_2, \psi_2)(T)$. We will show below that this implies $(A_1, \psi_1)(0) = (A_2, \psi_2)(0)$.

Define $(a_t, \varphi_t) = (A_2, \psi_2)(t) - (A_1, \psi_1)(t)$, $m_t = \mu_{(A_2, \psi_2)} - \mu_{(A_1, \psi_1)}$ and $n_t = \nu_{(A_2, \psi_2)} - \nu_{(A_1, \psi_1)}$. In terms of (a_t, φ_t) we can write

$$m_t = \mu_{(A_2, \psi_2)} - \mu_{(A_1, \psi_1)} = d_{A_1} a_t + a_t \wedge a_t + [\psi, \varphi_t] + \varphi_t \wedge \varphi_t$$

and for any $\eta \in \Omega^p(\text{End}(E))$ the difference of the associated Laplacians has the form

$$(A.7) \quad (\Delta_{(A_2, \psi_2)} - \Delta_{(A_1, \psi_1)}) \eta = \nabla_{A_1} a \times \eta + a \times \nabla_{A_1} \eta + a \times a \times \eta + \psi \times \varphi \times \eta + \varphi \times \varphi \times \eta$$

where again $\omega_1 \times \omega_2$ is used to denote a bilinear expression in ω_1 and ω_2 with constant coefficients. By definition of $\nu_{(A,\psi)}$ as the gradient of the Yang-Mills-Higgs functional at (d_A, ψ) we immediately have

$$\frac{\partial}{\partial t} (a_t + \varphi_t) = n_t, \quad \text{and} \quad \frac{\partial}{\partial t} (\nabla_{A_1} a_t + \nabla_{A_1} \varphi_t) = \left(\frac{\partial A}{\partial t} \times a_t, \frac{\partial A}{\partial t} \times \varphi_t \right) + \nabla_{A_1} n_t$$

Equation (A.5) then becomes

$$\begin{aligned} \left(\frac{\partial}{\partial t} + \Delta_{(A_1, \psi_1)} \right) m_t &= - (\Delta_{(A_2, \psi_2)} - \Delta_{(A_1, \psi_1)}) \mu_{(A_2, \psi_2)} \\ &\quad + a_t \times \psi_1 \times (F_{A_1} + \psi_1 \wedge \psi_1) + \nabla_{A_1} \varphi_t \times (F_{A_1} + \psi_1 \wedge \psi_1) \\ &\quad + \nabla_{A_1} \psi_1 \times m_t + \psi_1 \times n_t \end{aligned}$$

and equation (A.6) becomes

$$\begin{aligned} \left(\frac{\partial}{\partial t} + \Delta_{(A_1, \psi_1)} \right) n_t &= *[\nu_{(A_2, \psi_2)}, *\mu_{(A_2, \psi_2)}] - *[\nu_{(A_1, \psi_1)}, *\mu_{(A_1, \psi_1)}] - (\Delta_{(A_2, \psi_2)} - \Delta_{(A_1, \psi_1)}) \nu_{(A_2, \psi_2)} \\ &= *[n_t, *\mu_{(A_2, \psi_2)}] + *[\nu_{(A_1, \psi_1)}, *m_t] - (\Delta_{(A_2, \psi_2)} - \Delta_{(A_1, \psi_1)}) \nu_{(A_2, \psi_2)} \end{aligned}$$

Using (A.7) and the Weitzenböck formula (A.2), we then have the following inequalities. In the case where X is a compact Riemann surface then the estimates of [51, Sec. 2.2] show that all of the derivatives of the connection, the Higgs field and the curvature F_A are uniformly bounded along the flow and so the constant can be chosen uniformly on the interval $[0, T]$.

Lemma A.2. *For any pair of solutions $(d_{A_1}, \psi_1)(t)$ and $(d_{A_2}, \psi_2)(t)$ to the Yang-Mills-Higgs flow (A.4) there exists a positive constant C (possibly depending on t) such that the following inequalities hold*

$$(A.8) \quad \left| \left(\frac{\partial}{\partial t} + \nabla_{A_1}^* \nabla_{A_1} \right) m_t \right| \leq C (|a_t| + |\varphi_t| + |\nabla_{A_1} a_t| + |\nabla_{A_1} \varphi_t| + |m_t| + |n_t|)$$

$$(A.9) \quad \left| \left(\frac{\partial}{\partial t} + \nabla_{A_1}^* \nabla_{A_1} \right) n_t \right| \leq C (|a_t| + |\varphi_t| + |\nabla_{A_1} a_t| + |\nabla_{A_1} \varphi_t| + |m_t| + |n_t|)$$

$$(A.10) \quad \left| \frac{\partial}{\partial t} (a_t + \varphi_t) \right| = |n_t|$$

$$(A.11) \quad \left| \frac{\partial}{\partial t} (\nabla_A a_t + \nabla_A \varphi_t) \right| \leq C (|a_t| + |\varphi_t| + |\nabla_A n_t|)$$

Moreover, if X is a compact Riemann surface then the constant C can be chosen uniformly on any finite time interval $[0, T]$.

For simplicity of notation, in the following we use $\nabla := \nabla_{A_1}$ and $\square := \nabla_{A_1}^* \nabla_{A_1}$. Let $X := (m_t, n_t)$ and $Y := (a_t, \varphi_t, \nabla a_t, \nabla \varphi_t)$. The previous lemma implies that there exists a positive constant C such that the following inequalities hold

$$(A.12) \quad \begin{aligned} \left| \frac{\partial X}{\partial t} + \square X \right| &\leq C (|X| + |\nabla X| + |Y|) \\ \left| \frac{\partial Y}{\partial t} \right| &\leq C (|X| + |\nabla X| + |Y|) \end{aligned}$$

A general result of Kotschwar in [28, Thm 3] shows that any system satisfying (A.12) on the time interval $[0, T]$ for which $X(T) = 0, Y(T) = 0$, must also satisfy $X(t) = 0, Y(t) = 0$ for all $t \in [0, T]$. In the context of the Yang-Mills-Higgs flow (A.4), this gives us the proof of Proposition A.1.

REFERENCES

- [1] M. F. Atiyah and R. Bott. The Yang-Mills equations over Riemann surfaces. *Philos. Trans. Roy. Soc. London Ser. A*, 308(1505):523–615, 1983.
- [2] Indranil Biswas and Graeme Wilkin. Morse theory for the space of Higgs G -bundles. *Geom. Dedicata*, 149:189–203, 2010.
- [3] Jean-Pierre Bourguignon and H. Blaine Lawson, Jr. Stability and isolation phenomena for Yang-Mills fields. *Comm. Math. Phys.*, 79(2):189–230, 1981.
- [4] Insong Choe and George H. Hitching. Secant varieties and Hirschowitz bound on vector bundles over a curve. *Manuscripta Math.*, 133(3-4):465–477, 2010.
- [5] Georgios D. Daskalopoulos. The topology of the space of stable bundles on a compact Riemann surface. *J. Differential Geom.*, 36(3):699–746, 1992.
- [6] Georgios D. Daskalopoulos and Richard A. Wentworth. Convergence properties of the Yang-Mills flow on Kähler surfaces. *J. Reine Angew. Math.*, 575:69–99, 2004.
- [7] S. K. Donaldson. A new proof of a theorem of Narasimhan and Seshadri. *J. Differential Geom.*, 18(2):269–277, 1983.
- [8] S. K. Donaldson. Anti self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles. *Proc. London Math. Soc. (3)*, 50(1):1–26, 1985.
- [9] S. K. Donaldson. Infinite determinants, stable bundles and curvature. *Duke Math. J.*, 54(1):231–247, 1987.
- [10] V. G. Drinfel'd. Two-dimensional l -adic representations of the fundamental group of a curve over a finite field and automorphic forms on $GL(2)$. *Amer. J. Math.*, 105(1):85–114, 1983.
- [11] Andreas Floer. Morse theory for Lagrangian intersections. *J. Differential Geom.*, 28(3):513–547, 1988.
- [12] Andreas Floer. A relative Morse index for the symplectic action. *Comm. Pure Appl. Math.*, 41(4):393–407, 1988.
- [13] Andreas Floer. The unregularized gradient flow of the symplectic action. *Comm. Pure Appl. Math.*, 41(6):775–813, 1988.
- [14] Edward Frenkel. Lectures on the Langlands program and conformal field theory. In *Frontiers in number theory, physics, and geometry. II*, pages 387–533. Springer, Berlin, 2007.

- [15] N. J. Hitchin. The self-duality equations on a Riemann surface. *Proc. London Math. Soc. (3)*, 55(1):59–126, 1987.
- [16] N. J. Hitchin. Stable bundles and integrable systems. *Duke Math. J.*, 54(1):91–114, 1987.
- [17] George H. Hitching. Geometry of vector bundle extensions and applications to a generalised theta divisor. *Math. Scand.*, 112(1):61–77, 2013.
- [18] Min-Chun Hong. Heat flow for the Yang-Mills-Higgs field and the Hermitian Yang-Mills-Higgs metric. *Ann. Global Anal. Geom.*, 20(1):23–46, 2001.
- [19] John H. Hubbard. Parametrizing unstable and very unstable manifolds. *Mosc. Math. J.*, 5(1):105–124, 2005.
- [20] Daniel Huybrechts and Manfred Lehn. *The geometry of moduli spaces of sheaves*. Aspects of Mathematics, E31. Friedr. Vieweg & Sohn, Braunschweig, 1997.
- [21] Jun-Muk Hwang and S. Ramanan. Hecke curves and Hitchin discriminant. *Ann. Sci. École Norm. Sup. (4)*, 37(5):801–817, 2004.
- [22] Adam Jacob. The limit of the Yang-Mills flow on semi-stable bundles. *J. Reine Angew. Math.*, 709:1–13, 2015.
- [23] Rémi Janner and Jan Swoboda. Elliptic Yang-Mills flow theory. *Math. Nachr.*, 288(8-9):935–967, 2015.
- [24] Jürgen Jost. *Riemannian geometry and geometric analysis*. Universitext. Springer-Verlag, Berlin, fourth edition, 2005.
- [25] Anton Kapustin and Edward Witten. Electric-magnetic duality and the geometric Langlands program. *Commun. Number Theory Phys.*, 1(1):1–236, 2007.
- [26] Frances Clare Kirwan. *Cohomology of quotients in symplectic and algebraic geometry*, volume 31 of *Mathematical Notes*. Princeton University Press, Princeton, NJ, 1984.
- [27] Shoshichi Kobayashi. *Differential geometry of complex vector bundles*, volume 15 of *Publications of the Mathematical Society of Japan*. Princeton University Press, Princeton, NJ, 1987. , Kano Memorial Lectures, 5.
- [28] Brett Kotschwar. A short proof of backward uniqueness for some geometric evolution equations. *Internat. J. Math.*, 27(12):1650102, 17, 2016.
- [29] H. Lange and M. S. Narasimhan. Maximal subbundles of rank two vector bundles on curves. *Math. Ann.*, 266(1):55–72, 1983.
- [30] Herbert Lange. Zur Klassifikation von Regelmännigfaltigkeiten. *Math. Ann.*, 262(4):447–459, 1983.
- [31] Gérard Laumon. Correspondance de Langlands géométrique pour les corps de fonctions. *Duke Math. J.*, 54(2):309–359, 1987.
- [32] Jiayu Li and Xi Zhang. The gradient flow of Higgs pairs. *J. Eur. Math. Soc. (JEMS)*, 13(5):1373–1422, 2011.
- [33] S. Łojasiewicz. Sur les ensembles semi-analytiques. pages 237–241, 1971.
- [34] Hisashi Naito, Hideo Kozono, and Yoshiaki Maeda. A stable manifold theorem for the Yang-Mills gradient flow. *Tohoku Math. J. (2)*, 42(1):45–66, 1990.
- [35] M. S. Narasimhan and S. Ramanan. Moduli of vector bundles on a compact Riemann surface. *Ann. of Math. (2)*, 89:14–51, 1969.
- [36] M. S. Narasimhan and S. Ramanan. Vector bundles on curves. In *Algebraic Geometry (Internat. Colloq., Tata Inst. Fund. Res., Bombay, 1968)*, pages 335–346. Oxford Univ. Press, London, 1969.
- [37] M. S. Narasimhan and S. Ramanan. Deformations of the moduli space of vector bundles over an algebraic curve. *Ann. Math. (2)*, 101:391–417, 1975.
- [38] M. S. Narasimhan and S. Ramanan. Geometry of Hecke cycles. I. In *C. P. Ramanujam—a tribute*, volume 8 of *Tata Inst. Fund. Res. Studies in Math.*, pages 291–345. Springer, Berlin-New York, 1978.
- [39] M. S. Narasimhan and C. S. Seshadri. Stable and unitary vector bundles on a compact Riemann surface. *Ann. of Math. (2)*, 82:540–567, 1965.
- [40] Edward Nelson. *Topics in dynamics. I: Flows*. Mathematical Notes. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1969.
- [41] Johan Råde. On the Yang-Mills heat equation in two and three dimensions. *J. Reine Angew. Math.*, 431:123–163, 1992.
- [42] Michael Reed and Barry Simon. *Methods of modern mathematical physics. III. Scattering theory*. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1979.
- [43] Benjamin Sibley. Asymptotics of the Yang-Mills flow for holomorphic vector bundles over Kähler manifolds: the canonical structure of the limit. *J. Reine Angew. Math.*, 706:123–191, 2015.
- [44] Leon Simon. Asymptotics for a class of nonlinear evolution equations, with applications to geometric problems. *Ann. of Math. (2)*, 118(3):525–571, 1983.
- [45] Carlos T. Simpson. Constructing variations of Hodge structure using Yang-Mills theory and applications to uniformization. *J. Amer. Math. Soc.*, 1(4):867–918, 1988.
- [46] Carlos T. Simpson. Higgs bundles and local systems. *Inst. Hautes Études Sci. Publ. Math.*, (75):5–95, 1992.
- [47] Reyner Sjamaar. Holomorphic slices, symplectic reduction and multiplicities of representations. *Ann. of Math. (2)*, 141(1):87–129, 1995.
- [48] Jan Swoboda. Morse homology for the Yang-Mills gradient flow. *J. Math. Pures Appl. (9)*, 98(2):160–210, 2012.
- [49] K. Uhlenbeck and S.-T. Yau. On the existence of Hermitian-Yang-Mills connections in stable vector bundles. *Comm. Pure Appl. Math.*, 39(S, suppl.):S257–S293, 1986. *Frontiers of the mathematical sciences: 1985* (New York, 1985).

- [50] Joa Weber. A backward λ -lemma for the forward heat flow. *Math. Ann.*, 359(3-4):929–967, 2014.
- [51] Graeme Wilkin. Morse theory for the space of Higgs bundles. *Comm. Anal. Geom.*, 16(2):283–332, 2008.
- [52] Edward Witten. More on Gauge Theory and Geometric Langlands. *Adv. Math.*, 327:624–707, 2018.

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE 119076
E-mail address: graeme@nus.edu.sg